

INVOLUTIONS ON PRO- p -IWAHORI HECKE ALGEBRAS

NORIYUKI ABE

ABSTRACT. The pro- p -Iwahori Hecke algebra has an involution ι defined in terms of Iwahori-Matsumoto basis. Then for a module π of pro- p -Iwahori Hecke, $\pi^\iota = \pi \circ \iota$ is also a module. We calculate π^ι for simple modules π . We also calculate the dual of π .

1. INTRODUCTION

This is the sequel of [Abe16] and the aim of these papers are to calculate the extension of simple modules of pro- p -Iwahori Hecke algebras. The calculation will be appeared in sequel and we will use the results of this paper.

Let G be a connected reductive group over a non-archimedean local field with residue characteristic p . For a field C , we can attach the pro- p -Iwahori Hecke algebra of G . This is the convolution algebra of compactly supported functions which is bi-invariant under the pro- p radical of an Iwahori subgroup. If the characteristic of C is p , then this algebra plays an important role for the representation theory of G over C (cf. [AHHV17]).

The main object of this paper are the anti-involution ζ and the involution ι when the characteristic of C is p . These are defined as follows.

- ζ : Let $W(1)$ be the “pro- p Weyl group” (see subsection 2.2 for the precise definition). Then $\mathcal{H}$ has a basis $\{T_w \mid w \in W(1)\}$ parametrized by $W(1)$ which is called *Iwahori-Matsumoto basis*. The anti-involution ζ is defined by $\zeta(T_w) = T_{w^{-1}}$.
- ι : We also have another basis of $\mathcal{H}$ denoted by $\{T_w^* \mid w \in W(1)\}$. Then the involution ι is defined by $\iota(T_w) = (-1)^{\ell(w)} T_w^*$ where ℓ is the length function on $W(1)$.

By the multiplication rule of $\mathcal{H}$ in terms of the basis $\{T_w \mid w \in W(1)\}$ (the braid relations and the quadratic relations), these maps respect the multiplication.

Let π be a right $\mathcal{H}$ -module. Then we can attach the following two modules.

- $\pi^* = \text{Hom}_C(\pi, C)$ where the action of $X \in \mathcal{H}$ on $f \in \pi^*$ is given by $(fX)(v) = f(v\zeta(X))$ for $v \in \pi$.
- $\pi^\iota = \pi \circ \iota$.

If π is simple then π^* and π^ι are also simple. We determine these modules (Theorem 3.28, 4.9).

1.1. Organization of this paper. In the next section we give notation and recall some results on pro- p -Iwahori Hecke algebras. We use the same notation as [Abe16] and often refer this paper. In Section 3, we calculate π^ι for simple modules π . Since the construction of simple modules is divided into three steps, the calculation also has three steps, namely when π is the parabolic induction, generalized Steinberg

modules and supersingular modules. The hardest step is for the generalized Steinberg modules which will be calculated from subsection 3.2 to 3.7. The calculation of the dual of simple modules will be done in Section 4. In the calculation we use results in Section 3.

Acknowledgments. Most of this work was done during my pleasant stay at Institut de mathématiques de Jussieu. The work is supported by JSPS KAKENHI Grant Number 26707001.

2. PRELIMINARIES

2.1. Pro- p -Iwahori Hecke algebra. Let $\mathcal{H}$ be a pro- p -Iwahori Hecke algebra over a commutative ring C [Vig16]. We study modules over $\mathcal{H}$ in this paper. *In this paper, a module means a right module.* The algebra $\mathcal{H}$ is defined with a combinatorial data $(W_{\text{aff}}, S_{\text{aff}}, \Omega, W, W(1), Z_\kappa)$ and a parameter (q, c) .

We recall the definitions. The data satisfy the following.

- $(W_{\text{aff}}, S_{\text{aff}})$ is a Coxeter system.
- Ω acts on $(W_{\text{aff}}, S_{\text{aff}})$.
- $W = W_{\text{aff}} \rtimes \Omega$.
- Z_κ is a finite commutative group.
- The group $W(1)$ is an extension of W by Z_κ , namely we have an exact sequence $1 \rightarrow Z_\kappa \rightarrow W(1) \rightarrow W \rightarrow 1$.

The subgroup Z_κ is normal in $W(1)$. Hence the conjugate action of $w \in W(1)$ induces an automorphism of Z_κ , hence of the group ring $C[Z_\kappa]$. We denote it by $c \mapsto w \cdot c$.

Let $\text{Ref}(W_{\text{aff}})$ be the set of reflections in W_{aff} and $\text{Ref}(W_{\text{aff}}(1))$ the inverse image of $\text{Ref}(W_{\text{aff}})$ in $W(1)$. The parameter (q, c) maps $q: S_{\text{aff}} \rightarrow C$ and $c: \text{Ref}(W_{\text{aff}}(1)) \rightarrow C[Z_\kappa]$ with the following conditions. (Here the image of s by q (resp. c) is denoted by q_s (resp. c_s).

- For $w \in W$ and $s \in S_{\text{aff}}$, if $ws w^{-1} \in S_{\text{aff}}$ then $q_{ws w^{-1}} = q_s$.
- For $w \in W(1)$ and $s \in \text{Ref}(W_{\text{aff}}(1))$, $c_{ws w^{-1}} = w \cdot c_s$.
- For $s \in \text{Ref}(W_{\text{aff}}(1))$ and $t \in Z_\kappa$, we have $c_{ts} = t c_s$.

Let $S_{\text{aff}}(1)$ be the inverse image of S_{aff} in $W(1)$. For $s \in S_{\text{aff}}(1)$, we write q_s for $q_{\bar{s}}$ where $\bar{s} \in S_{\text{aff}}$ is the image of s . The length function on W_{aff} is denoted by ℓ and its inflation to W and $W(1)$ is also denoted by ℓ .

The C -algebra $\mathcal{H}$ is a free C -module and has a basis $\{T_w\}_{w \in W(1)}$. The multiplication is given by

- (Quadratic relations) $T_s^2 = q_s T_{s^2} + c_s T_s$ for $s \in S_{\text{aff}}(1)$.
- (Braid relations) $T_{vw} = T_v T_w$ if $\ell(vw) = \ell(v) + \ell(w)$.

We extend $q: S_{\text{aff}} \rightarrow C$ to $q: W \rightarrow C$ as follows. For $w \in W$, take $s_1, \dots, s_l \in S_{\text{aff}}$ and $u \in \Omega$ such that $w = s_1 \cdots s_l u$. Then put $q_w = q_{s_1} \cdots q_{s_l}$. From the definition, we have $q_{w^{-1}} = q_w$. We also put $q_w = q_{\bar{w}}$ for $w \in W(1)$ with the image $\bar{w}$ in W .

2.2. The data from a group. Let F be a non-archimedean local field, κ its residue field, p its residue characteristic and G a connected reductive group over F . We can get the data in the previous subsection from G as follows. See [Vig16], especially 3.9 and 4.2 for the details.

Fix a maximal split torus S and denote the centralizer of S by Z . Let Z^0 be the unique parahoric subgroup of Z and $Z(1)$ its pro- p radical. Then the group $W(1)$

(resp. W) is defined by $W(1) = N_G(Z)/Z(1)$ (resp. $W = N_G(Z)/Z^0$) where $N_G(Z)$ is the normalizer of Z in G . We also have $Z_\kappa = Z^0/Z(1)$. Let G' be the group generated by the unipotent radical of parabolic subgroups [AHHV17, II.1] and W_{aff} the image of $G' \cap N_G(Z)$ in W . Then this is a Coxeter group. Fix a set of simple reflections S_{aff} . The group W has the natural length function and let Ω be the set of length zero elements in W . Then we get the data $(W_{\text{aff}}, S_{\text{aff}}, \Omega, W, W(1), Z_\kappa)$.

Consider the apartment attached to S and an alcove surrounded by the hyperplanes fixed by S_{aff} . Let $I(1)$ be the pro- p -Iwahori subgroup attached to this alcove. Then with $q_s = \#(I(1)\tilde{s}I(1)/I(1))$ for $s \in S_{\text{aff}}$ with a lift $\tilde{s} \in N_G(Z)$ and suitable c_s , the algebra $\mathcal{H}$ is isomorphic to the Hecke algebra attached to $(G, I(1))$ [Vig16, Proposition 4.4].

In this paper, *the data $(W_{\text{aff}}, S_{\text{aff}}, \Omega, W, W(1), Z_\kappa)$ and the parameters (q, c) come from G in this way.* Let $W_{\text{aff}}(1)$ be the image of $G' \cap N_G(Z)$ in $W(1)$ and put $\mathcal{H}_{\text{aff}} = \bigoplus_{w \in W_{\text{aff}}(1)} CT_w$. This is a subalgebra of $\mathcal{H}$.

2.3. The algebra $\mathcal{H}[q_s]$ and $\mathcal{H}[q_s^{\pm 1}]$. For each $s \in S_{\text{aff}}$, let $\mathbf{q}_s$ be an indeterminate such that if $ws w^{-1} \in S_{\text{aff}}$ for $w \in W$, we have $\mathbf{q}_{ws w^{-1}} = \mathbf{q}_s$. Let $C[\mathbf{q}_s]$ be a polynomial ring with these indeterminate. Then with the parameter $s \mapsto \mathbf{q}_s$ and the other data coming from G , we have the algebra. This algebra is denoted by $\mathcal{H}[\mathbf{q}_s]$ and we put $\mathcal{H}[\mathbf{q}_s^{\pm 1}] = \mathcal{H}[\mathbf{q}_s] \otimes_{C[\mathbf{q}_s]} C[\mathbf{q}_s^{\pm 1}]$. Under $\mathbf{q}_s \mapsto \#(I(1)\tilde{s}I(1)/I(1)) \in C$ where $\tilde{s} \in N_G(Z)$ is a lift of s , we have $\mathcal{H}[\mathbf{q}_s] \otimes_{C[\mathbf{q}_s]} C \simeq \mathcal{H}$. As an abbreviation, we denote $\mathbf{q}_s$ by just q_s . Consequently we denote by $\mathcal{H}[q_s]$ (resp. $\mathcal{H}[q_s^{\pm 1}]$).

Since q_s is invertible in $\mathcal{H}[q_s^{\pm 1}]$, we can do some calculations in $\mathcal{H}[q_s^{\pm 1}]$ with q_s^{-1} . If the result can be stated in $\mathcal{H}[q_s]$, then this is an equality in $\mathcal{H}[q_s]$ since $\mathcal{H}[q_s]$ is a subalgebra of $\mathcal{H}[q_s^{\pm 1}]$ and by specializing, we can get some equality in $\mathcal{H}$. See [Vig16, 4.5] for more details.

2.4. The root system and the Weyl groups. Let $W_0 = N_G(Z)/Z$ be the finite Weyl group. Then this is a quotient of W . Recall that we have the alcove defining $I(1)$. Fix a special point $\mathbf{x}_0$ from the border of this alcove. Then $W_0 \simeq \text{Stab}_W \mathbf{x}_0$ and the inclusion $\text{Stab}_W \mathbf{x}_0 \hookrightarrow W$ is a splitting of the canonical projection $W \rightarrow W_0$. Throughout this paper, we fix this special point and regard W_0 as a subgroup of W . Set $S_0 = S_{\text{aff}} \cap W_0 \subset W$. This is a set of simple reflections in W_0 . For each $w \in W_0$, we fix a representative $n_w \in W(1)$ such that $n_{w_1 w_2} = n_{w_1} n_{w_2}$ if $\ell(w_1 w_2) = \ell(w_1) + \ell(w_2)$.

The group W_0 is the Weyl group of the root system Σ attached to (G, S) . Our fixed alcove and special point give a positive system of Σ , denoted by Σ^+ . The set of simple roots is denoted by Δ . As usual, for $\alpha \in \Delta$, let $s_\alpha \in S_0$ be a simple reflection for α .

The kernel of $W(1) \rightarrow W_0$ (resp. $W \rightarrow W_0$) is denoted by $\Lambda(1)$ (resp. Λ). Then $Z_\kappa \subset \Lambda(1)$ and we have $\Lambda = \Lambda(1)/Z_\kappa$. The group Λ (resp. $\Lambda(1)$) is isomorphic to Z/Z^0 (resp. $Z/Z(1)$). Any element in $W(1)$ can be uniquely written as $n_w \lambda$ where $w \in W_0$ and $\lambda \in \Lambda(1)$. We have $W = W_0 \ltimes \Lambda$.

2.5. The map ν . The group W acts on the apartment attached to S and the action of Λ is by the translation. Since the group of translations of the apartment is $X_*(S) \otimes_{\mathbb{Z}} \mathbb{R}$, we have a group homomorphism $\nu: \Lambda \rightarrow X_*(S) \otimes_{\mathbb{Z}} \mathbb{R}$. The compositions $\Lambda(1) \rightarrow \Lambda \rightarrow X_*(S) \otimes_{\mathbb{Z}} \mathbb{R}$ and $Z \rightarrow \Lambda \rightarrow X_*(S) \otimes_{\mathbb{Z}} \mathbb{R}$ are also denoted by ν . The homomorphism $\nu: Z \rightarrow X_*(S) \otimes_{\mathbb{Z}} \mathbb{R} \simeq \text{Hom}_{\mathbb{Z}}(X^*(S), \mathbb{R})$ is characterized by the following: For $t \in S$ and $\chi \in X^*(S)$, we have $\nu(t)(\chi) = -\text{val}(\chi(t))$ where val is

the normalized valuation of F . The kernel of $\nu: Z \rightarrow X_*(S) \otimes_{\mathbb{Z}} \mathbb{R}$ is equal to the maximal compact subgroup $\tilde{Z}$ of Z . In particular, $\text{Ker}(\Lambda(1) \xrightarrow{\nu} X_*(S) \otimes_{\mathbb{Z}} \mathbb{R}) = \tilde{Z}/Z(1)$ is a finite group.

We call $\lambda \in \Lambda(1)$ dominant (resp. anti-dominant) if $\nu(\lambda)$ is dominant (resp. anti-dominant).

Since the group W_{aff} is a Coxeter system, it has the Bruhat order denoted by $\leq$. For $w_1, w_2 \in W_{\text{aff}}$, we write $w_1 < w_2$ if there exists $u \in \Omega$ such that $w_1 u, w_2 u \in W_{\text{aff}}$ and $w_1 u < w_2 u$. Moreover, for $w_1, w_2 \in W(1)$, we write $w_1 < w_2$ if $w_1 \in W_{\text{aff}}(1)w_2$ and $\overline{w}_1 < \overline{w}_2$ where $\overline{w}_1, \overline{w}_2$ are the image of w_1, w_2 in W , respectively. We write $w_1 \leq w_2$ if $w_1 < w_2$ or $w_1 = w_2$.

2.6. Other basis. For $w \in W(1)$, take $s_1, \dots, s_l \in S_{\text{aff}}(1)$ and $u \in W(1)$ such that $l = \ell(w)$, $\ell(u) = 0$ and $w = s_1 \cdots s_l u$. Set $T_w^* = (T_{s_1} - c_{s_1}) \cdots (T_{s_l} - c_{s_l}) T_u$. Then this does not depend on the choice and we have $T_w^* \in T_w + \sum_{v < w} CT_v$. In particular, $\{T_w^*\}_{w \in W(1)}$ is a basis of $\mathcal{H}$. In $\mathcal{H}[q_s^{\pm 1}]$, we have $T_w^* = q_w T_{w^{-1}}^{-1}$.

For simplicity, we always assume that our commutative ring C contains a square root of q_s which is denoted by $q_s^{1/2}$ for $s \in S_{\text{aff}}$. For $w = s_1 \cdots s_l u$ where $\ell(w) = l$ and $\ell(u) = 0$, $q_w^{1/2} = q_{s_1}^{1/2} \cdots q_{s_l}^{1/2}$ is a square root of q_w . For a spherical orientation o , there is a basis $\{E_o(w)\}_{w \in W(1)}$ of $\mathcal{H}$ introduced in [Vig16, 5]. We have

$$E_o(w) \in T_w + \sum_{v < w} CT_v.$$

This satisfies the following product formula [Vig16, Theorem 5.25].

$$(2.1) \quad E_o(w_1) E_{o \cdot w_1}(w_2) = q_{w_1 w_2}^{-1/2} q_{w_1}^{1/2} q_{w_2}^{1/2} E_o(w_1 w_2).$$

Remark 2.1. Since we do not assume that q_s is invertible in C , $q_{w_1 w_2}^{-1/2} q_{w_1}^{1/2} q_{w_2}^{1/2}$ does not make sense in a usual way. See [Abe16, Remark 2.2].

2.7. Levi subalgebra. Since we have a positive system Σ^+ , we have a minimal parabolic subgroup B with a Levi part Z . In this paper, *parabolic subgroups are always standard, namely containing B* . Note that such parabolic subgroups correspond to subsets of Δ .

Let P be a parabolic subgroup. Attached to the Levi part of P containing Z , we have the data $(W_{\text{aff}, P}, S_{\text{aff}, P}, \Omega_P, W_P, W_P(1), Z_\kappa)$ and the parameters (q_P, c_P) . Hence we have the algebra $\mathcal{H}$. The parameter c_P is given by the restriction of c , hence we denote it just by c . The parameter q_P is defined as in [Abe, 4.1].

For the objects attached to this data, we add the suffix P . We have the set of simple roots Δ_P , the root system Σ_P and its positive system Σ_P^+ , the finite Weyl group $W_{0, P}$, the set of simple reflections $S_{0, P} \subset W_{0, P}$, the length function ℓ_P and the base $\{T_w^P\}_{w \in W_P(1)}$, $\{T_w^{P*}\}_{w \in W_P(1)}$ and $\{E_o^P(w)\}_{w \in W_P(1)}$ of $\mathcal{H}_P$. Note that we have no Λ_P , $\Lambda_P(1)$ and $Z_{\kappa, P}$ since they are equal to Λ , $\Lambda(1)$ and Z_κ .

An element $w = n_v \lambda \in W_P(1)$ where $v \in W_{0, P}$ and $\lambda \in \Lambda(1)$ is called P -positive (resp. P -negative) if $\langle \alpha, \nu(\lambda) \rangle \leq 0$ (resp. $\langle \alpha, \nu(\lambda) \rangle \geq 0$) for any $\alpha \in \Sigma^+ \setminus \Sigma_P^+$. Let $W_P^+(1)$ (resp. $W_P^-(1)$) be the set of P -positive (resp. P -negative) elements and put $\mathcal{H}_P^\pm = \bigoplus_{w \in W_P^\pm(1)} CT_w^P$. These are subalgebras of $\mathcal{H}_P$ [Abe, Lemma 4.1].

Proposition 2.2 ([Vig15b, Theorem 1.4]). *Let λ_P^+ (resp. λ_P^-) be in the center of $W_P(1)$ such that $\langle \alpha, \nu(\lambda_P^+) \rangle < 0$ (resp. $\langle \alpha, \nu(\lambda_P^-) \rangle > 0$) for all $\alpha \in \Sigma^+ \setminus \Sigma_P^+$. Then*

$T_{\lambda_P^+}^P = T_{\lambda_P^+}^{P*} = E_{o_-,P}^P(\lambda_P^+)$ (resp. $T_{\lambda_P^-}^P = T_{\lambda_P^-}^{P*} = E_{o_-,P}^P(\lambda_P^-)$) is in the center of $\mathcal{H}_P$ and we have $\mathcal{H}_P = \mathcal{H}_P^+ E_{o_-,P}^P(\lambda_P^+)^{-1}$ (resp. $\mathcal{H}_P = \mathcal{H}_P^- E_{o_-,P}^P(\lambda_P^-)^{-1}$).

Note that such $\lambda_P^\pm$ always exists [Abe16, Lemma 2.4].

We define $j_P^\pm: \mathcal{H}_P^\pm \rightarrow \mathcal{H}$ and $j_P^{\pm*}: \mathcal{H}_P^\pm \rightarrow \mathcal{H}$ by $j_P^\pm(T_w^P) = T_w$ and $j_P^{\pm*}(T_w^{P*}) = T_w^*$ for $w \in W_P^\pm(1)$. Then these are algebra homomorphisms.

Let Q be a parabolic subgroup containing P and let $W_P^{Q+}(1)$ (resp. $W_P^{Q-}(1)$) be the set of $n_w \lambda$ where $\langle \alpha, \nu(\lambda) \rangle \leq 0$ (resp. $\langle \alpha, \nu(\lambda) \rangle \geq 0$) for any $\alpha \in \Sigma_Q^+ \setminus \Sigma_P^+$ and $w \in W_{0,P}$. Put $\mathcal{H}_P^{Q\pm} = \bigoplus_{w \in W_P^{Q\pm}(1)} CT_w^P \subset \mathcal{H}_P$. Then we have homomorphisms $j_P^{Q\pm}, j_P^{Q\pm*}: \mathcal{H}_P^{Q\pm} \rightarrow \mathcal{H}_Q$ defined by a similar way.

2.8. Parabolic induction. Let P be the parabolic subgroup and σ an $\mathcal{H}_P$ -module. (This is a right module as in subsection 2.1.) Then we define an $\mathcal{H}$ -module $I_P(\sigma)$ by

$$I_P(\sigma) = \text{Hom}_{(\mathcal{H}_P^-, j_P^{-*})}(\mathcal{H}, \sigma).$$

We call I_P the parabolic induction. For $P \subset P_1$, we write

$$I_P^{P_1}(\sigma) = \text{Hom}_{(\mathcal{H}_P^{P_1-}, j_P^{P_1-*})}(\mathcal{H}_{P_1}, \sigma).$$

Let P be a parabolic subgroup. Set $W_0^P = \{w \in W_0 \mid w(\Delta_P) \subset \Sigma^+\}$. Then the multiplication map $W_0^P \times W_{0,P} \rightarrow W_0$ is bijective and for $w_1 \in W_0^P$ and $w_2 \in W_{0,P}$, we have $\ell(w_1 w_2) = \ell(w_1) + \ell(w_2)$. We also put ${}^P W_0 = \{w \in W_0 \mid w^{-1}(\Delta_P) \subset \Sigma^+\}$. Then the multiplication map $W_{0,P} \times {}^P W_0 \rightarrow W_0$ is bijective and for $w_1 \in W_{0,P}$ and $w_2 \in {}^P W_0$, we have $\ell(w_1 w_2) = \ell(w_1) + \ell(w_2)$. See [Abe16, Proposition 2.9] for the following proposition.

Proposition 2.3. *Let P be a parabolic subgroup and σ an $\mathcal{H}_P$ -module.*

- (1) *The map $I_P(\sigma) \ni \varphi \mapsto (\varphi(T_{n_w}))_{w \in W_0^P} \in \bigoplus_{w \in W_0^P} \sigma$ is bijective.*
- (2) *The map $I_P(\sigma) \ni \varphi \mapsto (\varphi(T_{n_w}^*))_{w \in W_0^P} \in \bigoplus_{w \in W_0^P} \sigma$ is bijective.*

Proposition 2.4 ([Abe, Proposition 4.12]). *Assume that $q_s = 0$ for any $s \in S_{\text{aff}}$. Let $w \in W_0^P$ and $\lambda \in \Lambda(1)$. Then for $\varphi \in I_P(\sigma)$, we have*

$$(\varphi E_{o_-}(\lambda))(T_{n_w}) = \begin{cases} \varphi(T_{n_w}) \sigma(E_{o_-,P}^P(n_w^{-1} \cdot \lambda)) & (n_w^{-1} \cdot \lambda \in W_P^-(1)), \\ 0 & (n_w^{-1} \cdot \lambda \notin W_P^-(1)). \end{cases}$$

We also define

$$I_P'(\sigma) = \text{Hom}_{(\mathcal{H}_P^-, j_P^-)}(\mathcal{H}, \sigma)$$

and $I_P^{P_1'}$ by the similar way.

2.9. Twist by $n_{w_G w_P}$. For a parabolic subgroup P , let w_P be the longest element in $W_{0,P}$. In particular, w_G is the longest element in W_0 . Let P' be a parabolic subgroup corresponding to $-w_G(\Delta_P)$, in other words, $P' = n_{w_G w_P} P^{\text{op}} n_{w_G w_P}^{-1}$ where P^{op} is the opposite parabolic subgroup of P with respect to the Levi part of P containing Z . Set $n = n_{w_G w_P}$. Then the map $P^{\text{op}} \rightarrow P'$ defined by $p \mapsto npn^{-1}$ is an isomorphism which preserves the data used to define the pro- p -Iwahori Hecke algebras. Hence $T_w^P \mapsto T_{nwn^{-1}}^{P'}$ gives an isomorphism $\mathcal{H}_P \rightarrow \mathcal{H}_{P'}$. This sends T_w^{P*} to $T_{nwn^{-1}}^{P'*}$ and $E_{o_+,P}^P(w)$ to $E_{o_+,P'}^{P'}(nwn^{-1})$ where $v \in W_{0,P}$.

Let σ be an $\mathcal{H}_P$ -module. Then we define an $\mathcal{H}_{P'}$ -module $n_{w_G w_P} \sigma$ via the pull-back of the above isomorphism. Namely, for $w \in W_{P'}(1)$, we put $(n_{w_G w_P} \sigma)(T_w^{P'}) = \sigma(T_{n_{w_G w_P}^{-1} w n_{w_G w_P}}^P)$.

2.10. The extension and the generalized Steinberg modules. Let P be a parabolic subgroup and σ an $\mathcal{H}_P$ -module. For $\alpha \in \Delta$, let P_α be a parabolic subgroup corresponding to $\Delta_P \cup \{\alpha\}$. Then we define $\Delta(\sigma) \subset \Delta$ by

$$\begin{aligned} \Delta(\sigma) &= \{\alpha \in \Delta \mid \langle \Delta_P, \alpha^\vee \rangle = 0, \sigma(T_\lambda^P) = 1 \text{ for any } \lambda \in W_{\text{aff}, P_\alpha}(1) \cap \Lambda(1)\} \cup \Delta_P. \end{aligned}$$

Let $P(\sigma)$ be a parabolic subgroup corresponding to $\Delta(\sigma)$.

Proposition 2.5 ([AHV17, Corollary 3.9]). *Let σ be an $\mathcal{H}_P$ -module and Q a parabolic subgroup between P and $P(\sigma)$. Denote the parabolic subgroup corresponding to $\Delta_Q \setminus \Delta_P$ by P_2 . Then there exist a unique $\mathcal{H}_Q$ -module $e_Q(\sigma)$ acting on the same space as σ such that*

- $e_Q(\sigma)(T_w^{Q*}) = \sigma(T_w^{P*})$ for any $w \in W_P(1)$.
- $e_Q(\sigma)(T_w^{Q*}) = 1$ for any $w \in W_{P_2, \text{aff}}(1)$.

Moreover, one of the following condition gives a characterization of $e_Q(\sigma)$.

- (1) For any $w \in W_P^{Q-}(1)$, $e_Q(\sigma)(T_w^{Q*}) = \sigma(T_w^{P*})$ (namely, $e_Q(\sigma) \simeq \sigma$ as $(\mathcal{H}_P^{Q-}, j_P^{Q-*})$ -modules) and for any $w \in W_{\text{aff}, P_2}(1)$, $e_Q(\sigma)(T_w^{Q*}) = 1$.
- (2) For any $w \in W_P^{Q+}(1)$, $e_Q(\sigma)(T_w^{Q*}) = \sigma(T_w^{P*})$ and for any $w \in W_{\text{aff}, P_2}(1)$, $e_Q(\sigma)(T_w^{Q*}) = 1$.

We call $e_Q(\sigma)$ the extension of σ to $\mathcal{H}_Q$. A typical example of the extension is the trivial representation $\mathbf{1} = \mathbf{1}_G$. This is a one-dimensional $\mathcal{H}$ -module defined by $\mathbf{1}(T_w) = q_w$, or equivalently $\mathbf{1}(T_w^*) = 1$. We have $\Delta(\mathbf{1}_P) = \{\alpha \in \Delta \mid \langle \Delta_P, \alpha^\vee \rangle = 0\} \cup \Delta_P$ and, if Q is a parabolic subgroup between P and $P(\mathbf{1}_P)$, we have $e_Q(\mathbf{1}_P) = \mathbf{1}_Q$.

Let $P(\sigma) \supset P_0 \supset Q_1 \supset Q \supset P$. Then as in [Abe, 4.5], we have $I_{Q_1}^{P_0}(e_{Q_1}(\sigma)) \subset I_Q^{P_0}(e_Q(\sigma))$. Define

$$\text{St}_Q^{P_0}(\sigma) = \text{Cok} \left(\bigoplus_{Q_1 \supsetneq Q} I_{Q_1}^{P_0}(e_{Q_1}(\sigma)) \rightarrow I_Q^{P_0}(e_Q(\sigma)) \right).$$

When $P_0 = G$, we write $\text{St}_Q(\sigma)$.

In the rest of this subsection, we assume that $P(\sigma) = G$. As we mentioned in the above, for $Q_1 \supset Q \supset P$, we have $I_{Q_1}(e_{Q_1}(\sigma)) \hookrightarrow I_Q(e_Q(\sigma))$. The proof of [Abe, Lemma 4.23] implies the following lemma.

Lemma 2.6. *Assume that $q_s = 0$ for any $s \in S_{\text{aff}}$. The diagram*

$$\begin{array}{ccc} I_{Q_1}(e_{Q_1}(\sigma)) & \hookrightarrow & I_Q(e_Q(\sigma)) \\ \wr \downarrow & & \downarrow \wr \\ \bigoplus_{w \in W_0^{Q_1}} \sigma & \hookrightarrow & \bigoplus_{w \in W_0^Q} \sigma \end{array}$$

is commutative. Here the embedding $\bigoplus_{w \in W_0^{Q_1}} \sigma \hookrightarrow \bigoplus_{w \in W_0^Q} \sigma$ is induced by $W_0^{Q_1} \hookrightarrow W_0^Q$.

Lemma 2.7. *Assume that $q_s = 0$ for any $s \in S_{\text{aff}}$. Let $\varphi \in I_Q(e_Q(\sigma))$. Then for $w \in W_0^Q$ and $\lambda \in \Lambda(1)$, we have*

$$(\varphi E_{o_-}(\lambda))(T_{n_w}) = \begin{cases} \varphi(T_{n_w})\sigma(E_{o_-,P}^P(n_w^{-1} \cdot \lambda)) & (n_w^{-1} \cdot \lambda \text{ is } P\text{-negative}), \\ 0 & (\text{otherwise}). \end{cases}$$

Proof. Assume that $n_w^{-1} \cdot \lambda$ is P -negative. Then in particular it is Q -negative. By Proposition 2.4, we have

$$(\varphi E_{o_-}(\lambda))(T_{n_w}) = \varphi(T_{n_w})e_Q(\sigma)(E_{o_-,Q}^Q(n_w^{-1} \cdot \lambda)).$$

Since $n_w^{-1} \cdot \lambda$ is P -negative, we have $E_{o_-,Q}^Q(n_w^{-1} \cdot \lambda) \in \mathcal{H}_P^{Q-}$. Hence $E_{o_-,Q}^Q(n_w^{-1} \cdot \lambda) = j_P^{Q-*}(E_{o_-,P}^P(n_w^{-1} \cdot \lambda))$ by [Abe16, Lemma 2.6]. Therefore we have $e_Q(\sigma)(E_{o_-,Q}^Q(n_w^{-1} \cdot \lambda)) = \sigma(E_{o_-,P}^P(n_w^{-1} \cdot \lambda))$. We get the lemma in this case.

Assume that $n_w^{-1} \cdot \lambda$ is not P -negative. Then there exists $\alpha \in \Sigma^+ \setminus \Sigma_P^+$ such that $\langle w(\alpha), \nu(\lambda) \rangle < 0$. Take λ_P^- as in Proposition 2.2 and put $\lambda_0 = n_w \cdot \lambda_P^-$. We have $\langle w(\alpha), \nu(\lambda_0) \rangle = \langle \alpha, \nu(\lambda_P^-) \rangle > 0$. Hence λ and λ_0 are not in the same chamber. Therefore we have $E_{o_-}(\lambda)E_{o_-}(\lambda_0) = 0$ by (2.1) and [Abe16, Lemma 2.11]. Hence we have $(\varphi E_{o_-}(\lambda)E_{o_-}(\lambda_0))(T_{n_w}) = 0$. Since $n_w^{-1} \cdot \lambda_0 = \lambda_P^-$ is P -negative, we have $(\varphi E_{o_-}(\lambda)E_{o_-}(\lambda_0))(T_{n_w}) = (\varphi E_{o_-}(\lambda))(T_{n_w})\sigma(E_{o_-,P}^P(\lambda_P^-))$ as we have already proved. Since $\lambda_P^- \in Z(W_P(1))$, $E_{o_-,P}^P(\lambda_P^-)$ is invertible. Hence we have $(\varphi E_{o_-}(\lambda))(T_{n_w}) = 0$. $\square$

2.11. Module $\sigma_{\ell-\ell_P}$. Let P be a parabolic subgroup and σ an $\mathcal{H}_P$ -module. Define a linear map $\sigma_{\ell-\ell_P}$ by $\sigma_{\ell-\ell_P}(T_w) = (-1)^{\ell(w)-\ell_P(w)}\sigma(T_w)$. Then this defines a new $\mathcal{H}_P$ -module $\sigma_{\ell-\ell_P}$ [Abe16, Lemma 4.1].

2.12. Supersingular modules. Assume that $q_s = 0$ for any $s \in S_{\text{aff}}$. Let $\mathcal{O}$ be a conjugacy class in $W(1)$ which is contained in $\Lambda(1)$. For a spherical orientation o , set $z_{\mathcal{O}} = \sum_{\lambda \in \mathcal{O}} E_o(\lambda)$. Then this does not depend on o and gives an element of the center of $\mathcal{H}$ [Vig15a, Theorem 5.1]. The length of $\lambda \in \mathcal{O}$ does not depend on λ . We denote it by $\ell(\mathcal{O})$.

Definition 2.8. Let π be an $\mathcal{H}$ -module. We call π supersingular if there exists $n \in \mathbb{Z}_{>0}$ such that $\pi z_{\mathcal{O}}^n = 0$ for any $\mathcal{O}$ such that $\ell(\mathcal{O}) > 0$.

The simple supersingular $\mathcal{H}$ -modules are classified in [Oll14, Vig15a]. We recall their results. Assume that C is a field. Let χ be a character of $Z_{\kappa} \cap W_{\text{aff}}(1)$ and put $S_{\text{aff},\chi} = \{s \in S_{\text{aff}} \mid \chi(c_s^-) \neq 0\}$ where $\tilde{s} \in W(1)$ is a lift of $s \in S_{\text{aff}}$. Note that if $\tilde{s}'$ is another lift, then $\tilde{s}' = t\tilde{s}$ for some $t \in Z_{\kappa}$. Hence $\chi(c_{\tilde{s}'}^-) = \chi(t)\chi(c_{\tilde{s}}^-)$. Therefore the condition does not depend on a choice of a lift. Let $J \subset S_{\text{aff},\chi}$. Then the character $\Xi = \Xi_{J,\chi}$ of $\mathcal{H}_{\text{aff}}$ is defined by

$$\begin{aligned} \Xi_{J,\chi}(T_t) &= \chi(t) \quad (t \in Z_{\kappa} \cap W_{\text{aff}}(1)), \\ \Xi_{J,\chi}(T_{\tilde{s}}^-) &= \begin{cases} \chi(c_{\tilde{s}}^-) & (s \in S_{\text{aff},\chi} \setminus J), \\ 0 & (s \notin S_{\text{aff},\chi} \setminus J) \end{cases} \end{aligned}$$

where $\tilde{s} \in W_{\text{aff}}(1)$ is a lift of s . Let $\Omega(1)_{\Xi}$ be the stabilizer of Ξ and V an simple $C[\Omega(1)_{\Xi}]$ -module such that $V|_{Z_{\kappa} \cap W_{\text{aff}}(1)}$ is a direct sum of χ . Put $\mathcal{H}_{\Xi} = \mathcal{H}_{\text{aff}}C[\Omega(1)_{\Xi}]$. This is a subalgebra of $\mathcal{H}$. For $X \in \mathcal{H}_{\text{aff}}$ and $Y \in C[\Omega(1)_{\Xi}]$, we define the action of XY on $\Xi \otimes V$ by $x \otimes y \mapsto xX \otimes yY$. Then this defines a well-defined action of $\mathcal{H}_{\Xi}$ on $\Xi \otimes V$. Set $\pi_{\chi,J,V} = (\Xi \otimes V) \otimes_{\mathcal{H}_{\Xi}} \mathcal{H}$.

Proposition 2.9 ([Vig15a, Theorem 1.6]). *The module $\pi_{\chi,J,V}$ is simple and it is supersingular if and only if the groups generated by J and generated by $S_{\text{aff},\chi} \setminus J$ are both finite. If C is an algebraically closed field, then any simple supersingular modules are given in this way.*

Remark 2.10. This classification result is valid even though the data which defines $\mathcal{H}$ does not come from a reductive group.

2.13. Simple modules. Assume that C is an algebraically closed field of characteristic p . We consider the following triple (P, σ, Q) .

- P is a parabolic subgroup.
- σ is an simple supersingular $\mathcal{H}_P$ -module.
- Q is a parabolic subgroup between P and $P(\sigma)$.

Define

$$I(P, \sigma, Q) = I_{P(\sigma)}(\text{St}_Q^{P(\sigma)}(\sigma)).$$

Theorem 2.11 ([Abe, Theorem 1.1]). *The module $I(P, \sigma, Q)$ is simple and any simple module has this form. Moreover, (P, σ, Q) is unique up to isomorphism.*

Let χ be a character of $Z_\kappa \cap W_{\text{aff},P}(1)$, $J \subset S_{\text{aff},P,\chi}$ and V an simple module of $C[\Omega_P(1)_{\Xi_{J,\chi}}]$ whose restriction to $Z_\kappa \cap W_{\text{aff},P}(1)$ is a direct sum of χ . Assume that the group generated by J and generated by $S_{P,\text{aff},\chi} \setminus J$ are finite. Then we put $I(P; \chi, J, V; Q) = I(P, \pi_{\chi,J,V}, Q)$. This is a simple module.

2.14. Assumption on C . In the rest of this paper, we always assume that $p = 0$ in C unless otherwise stated since almost all results in this paper is proved only under this assumption. Since q_s is a power of p , this assumption implies $q_w = 0$ for any $w \in W(1)$ such that $\ell(w) > 0$. When we discuss about simple modules, we also assume that C is a field. Such assumptions are written at the top of subsections or the statement of theorems.

3. TWIST

We define an involution $\iota = \iota_G: \mathcal{H} \rightarrow \mathcal{H}$ by $\iota(T_w) = (-1)^{\ell(w)} T_w^*$ [Vig16, Proposition 4.23] and $\pi^\iota = \pi \circ \iota$ for an $\mathcal{H}$ -module π . Obviously, π^ι is simple if π is simple. In this section, we calculate π^ι for simple modules π .

3.1. Parabolic induction. Let P be a parabolic subgroup. Then by [Abe16, Lemma 4.2], we have $(\sigma^{\iota P})_{\ell-\ell_P} = (\sigma_{\ell-\ell_P})^{\iota P}$. We denote this module by $\sigma_{\ell-\ell_P}^{\iota P}$. We have $I_P(\sigma)^\iota \simeq I'_P(\sigma_{\ell-\ell_P}^{\iota P})$ [Abe16, Proposition 4.11]. In this subsection, we prove the following proposition.

Proposition 3.1. *There exists a homomorphism $\Phi: I_P \rightarrow I'_P$ which is characterized by $\Phi(\varphi)(XT_{n_{w_G w_P}}) = \varphi(XT_{n_{w_G w_P}})$ for any $X \in \mathcal{H}$ and $\varphi \in I_P$.*

Remark 3.2. Assume that an $\mathcal{H}_P$ -module σ is not zero and $\varphi \in I_P(\sigma)$. Then we have $\Phi(\varphi)(T_{n_{w_G w_P}}) = \varphi(T_{n_{w_G w_P}})$. Since $I_P(\sigma) \rightarrow \sigma$ defined by $\varphi \mapsto \varphi(T_{n_{w_G w_P}})$ is surjective by Proposition 2.3, there exists $\varphi \in I_P(\sigma)$ such that $\varphi(T_{n_{w_G w_P}}) \neq 0$. Hence $\Phi(\varphi) \neq 0$. Therefore if σ is not zero, then $\Phi \neq 0$.

The following corollary is the first step to calculate π^ι for a simple $\mathcal{H}$ -module π .

Corollary 3.3. *Assume that C is a field. Let σ be an $\mathcal{H}_P$ -module. The representation $I_P(\sigma)$ is simple if and only if $I'_P(\sigma)$ is simple. Moreover, if it is the case, then $I_P(\sigma) \simeq I'_P(\sigma)$. Therefore if $I_P(\sigma)$ is simple, then $I_P(\sigma)^\iota \simeq I_P(\sigma_{\ell-P}^\iota)$.*

Proof. If $I_P(\sigma)$ is simple, then the homomorphism in Proposition 3.1 is injective. We prove that $\dim I_P(\sigma) = \dim I'_P(\sigma) < \infty$. Since $I_P(\sigma)$ is simple, σ is also simple. Hence it is finite-dimensional. By $I_P(\sigma) \simeq \bigoplus_{w \in W_0^P} \sigma$ (Proposition 2.3), we have $\dim I_P(\sigma) = \#W_0^P \dim \sigma$. We also have $\dim I'_P(\sigma) = \#W_0^P \dim \sigma$ by [Abe16, Proposition 4.12]. $\square$

For the proof of Proposition 3.1, by [Abe16, Proposition 4.13], it is sufficient to prove the following lemma.

Lemma 3.4. *Put $P' = n_{w_G w_P} P^{\text{op}} n_{w_G w_P}^{-1}$. The map $\varphi \mapsto (X \mapsto \varphi(X T_{n_{w_G w_P}}))$ gives a homomorphism*

$$I_P(\sigma) \rightarrow \text{Hom}_{(\mathcal{H}_{P'}, j_{P'}^+)}(\mathcal{H}, n_{w_G w_P} \sigma).$$

Proof. Set $n = n_{w_G w_P}$ and we prove $I_P(\sigma) \ni \varphi \mapsto \varphi(T_n) \in n\sigma$ is $(\mathcal{H}_{P'}, j_{P'}^+)$ -module homomorphism. Let $w \in W_{P'}(1)$ be a P' -positive element. By (4.1) in [Abe16], we have $j_{P'}^+(E_{o_+, P'}^{P'}(w))T_n = T_n j_P^-(E_{o_+, P}^P(n^{-1}wn))$. Hence

$$\begin{aligned} (\varphi j_{P'}^+(E_{o_+, P'}^{P'}(w)))(T_n) &= \varphi(j_{P'}^+(E_{o_+, P'}^{P'}(w))T_n) \\ &= \varphi(T_n j_P^-(E_{o_+, P}^P(n^{-1}wn))). \end{aligned}$$

We prove the following claim. From this claim, $(\varphi j_{P'}^+(E_{o_+, P'}^{P'}(w)))(T_n)$ only depends on $\varphi(T_n)$ and w .

Claim. Let $\varphi \in I_P(\sigma)$ and $X \in \mathcal{H}$. Then $\varphi(T_n X)$ only depends on $\varphi(T_n)$ and X .

We introduce a basis defined by

$$E_-(n_w \lambda) = q_{n_w \lambda}^{1/2} q_{n_w}^{-1/2} q_\lambda^{-1/2} T_{n_w}^* E_{o_-}(\lambda)$$

for $w \in W_0$ and $\lambda \in \Lambda(1)$. By [Abe, Lemma 4.2], $\{E_-(w) \mid w \in W(1)\}$ is a C -basis of $\mathcal{H}$.

To prove the claim, we may assume $X = E_-(n_w \lambda)$ for $w \in W_0$ and $\lambda \in \Lambda(1)$. Take $\lambda_P^- \in \Lambda(1)$ as in Proposition 2.2 such that $\lambda \lambda_P^-$ is P -negative. Then we have

$$\varphi(T_n E_-(n_w \lambda)) = \varphi(T_n E_-(n_w \lambda) E_{o_-}(\lambda_P^-)) \sigma(E_{o_-, P}^P(\lambda_P^-))^{-1}$$

by [Abe16, Lemma 2.6]. If $\ell(n_w \lambda) + \ell(\lambda_P^-) > \ell(n_w \lambda \lambda_P^-)$, then $E_-(n_w \lambda) E_{o_-}(\lambda_P^-) = 0$. Hence we have $\varphi(T_n E_-(n_w \lambda)) = 0$, so we get the claim. If $\ell(n_w \lambda) + \ell(\lambda_P^-) = \ell(n_w \lambda \lambda_P^-)$, then $E_-(n_w \lambda) E_{o_-}(\lambda_P^-) = E_-(n_w \lambda \lambda_P^-)$. Let $w_1 \in W_0^P$ and $w_2 \in W_{0, P}$ such that $w = w_1 w_2$. Then $n_{w_2} \lambda \lambda_P^- \in W_P(1)$ is P -negative. Hence $\ell(n_{w_1} n_{w_2} \lambda \lambda_P^-) = \ell(n_{w_1}) + \ell(n_{w_2} \lambda \lambda_P^-)$ by [Abe16, Lemma 2.18]. Therefore we have $E_-(n_w \lambda \lambda_P^-) = T_{n_{w_1}}^* E_-(n_{w_2} \lambda \lambda_P^-)$. If $w_1 \neq 1$, then $w_1 \notin W_{0, P}$. Hence in a reduced expression of w_1 , a simple reflection s_α for some $\alpha \in \Delta \setminus \Delta_P$ appears. Therefore, there exists $x \in W_{0, P}$, $\alpha \in \Delta \setminus \Delta_P$ and $y \in W_0$ such that $w_1 = x s_\alpha y$ and $\ell(w_1) = \ell(x) + \ell(s_\alpha) + \ell(y)$. Since $x \in W_{0, P}$, $x(\alpha) \in \Sigma^+ \setminus \Sigma_P^+$. Hence $w_P x(\alpha) > 0$.

Therefore $w_G w_P x(\alpha) < 0$. Hence $nn_x n_{s_\alpha} < nn_x$. Therefore we have

$$\begin{aligned} \ell(n) + \ell(w_1) &= \ell(w_G w_P) + \ell(x) + \ell(s_\alpha) + \ell(y) \\ &\geq \ell(w_G w_P x) + \ell(s_\alpha) + \ell(y) \\ &> \ell(w_G w_P x s_\alpha) + \ell(y) \\ &\geq \ell(w_G w_P x s_\alpha y) = \ell(nn_{w_1}). \end{aligned}$$

Hence $T_n T_{n_{w_1}}^* = E_{o_+, n^{-1}}(n) E_{o_+}(n_{w_1}) = 0$ by [Vig16, Example 5.22] and (2.1). Therefore, if $w_1 \neq 1$, namely $w \notin W_{0,P}$, then we have $\varphi(T_n E_-(n_w \lambda \lambda_P^-)) = \varphi(T_n T_{n_{w_1}}^* E_-(n_{w_2} \lambda \lambda_P^-)) = 0$ again. If $w \in W_{0,P}$, then $n_w \lambda \lambda_P^-$ is a P -negative element. Hence $E_-^P(n_w \lambda \lambda_P^-) \in \mathcal{H}_P^-$ and we have $E_-(n_w \lambda \lambda_P^-) = j_P^{-*}(E_-^P(n_w \lambda \lambda_P^-))$ by [Abe, Lemma 4.6]. Therefore we have

$$\begin{aligned} \varphi(T_n E_-(n_w \lambda)) &= \varphi(T_n E_-(n_w \lambda \lambda_P^-)) \sigma(E_{o_-, P}^P(\lambda_P^-))^{-1} \\ &= \varphi(T_n) \sigma(E_-^P(n_w \lambda \lambda_P^-) E_{o_-, P}^P(\lambda_P^-)^{-1}) \\ &= \varphi(T_n) \sigma(E_-^P(n_w \lambda)). \end{aligned}$$

The claim is proved.

Let $\varphi_0 \in I_P(\sigma)$ be such that $\varphi_0(T_n) = \varphi(T_n)$ and $\varphi_0(T_{n_v}) = 0$ for $v \in W_0^P \setminus \{w_G w_P\}$. Then, as a consequence of the claim, we have

$$(\varphi j_{P'}^+(E_{o_+, P'}^{P'}(w)))(T_n) = (\varphi_0 j_{P'}^+(E_{o_+, P'}^{P'}(w)))(T_n).$$

By the proof of [Abe, Proposition 4.14], we have

$$(\varphi_0 j_{P'}^+(E_{o_+, P'}^{P'}(w)))(T_n) = \varphi_0(T_n)(n\sigma)(E_{o_+, P'}^{P'}(w)).$$

Since $\varphi_0(T_n) = \varphi(T_n)$, we have

$$\varphi_0(T_n)(n\sigma)(E_{o_+, P'}^{P'}(w)) = \varphi(T_n)(n\sigma)(E_{o_+, P'}^{P'}(w)).$$

Hence

$$(\varphi j_{P'}^+(E_{o_+, P'}^{P'}(w)))(T_n) = \varphi(T_n)(n\sigma)(E_{o_+, P'}^{P'}(w)).$$

We get the lemma. $\square$

Here is the compatibility with the transitivity of I_P and I_P' [Abe16, Proposition 4.12].

Lemma 3.5. *Let $Q \supset P$ be a parabolic subgroups. Then the following three maps are equal.*

- (1) $I_P \rightarrow I_P'$.
- (2) $I_P = I_Q \circ I_P^Q \rightarrow I_Q' \circ I_P^Q \rightarrow I_Q' \circ I_P^{Q'} = I_P'$.
- (3) $I_P = I_Q \circ I_P^Q \rightarrow I_Q \circ I_P^{Q'} \rightarrow I_Q' \circ I_P^{Q'} = I_P'$.

Proof. Let σ be an $\mathcal{H}_P$ -module. In each cases, let $\varphi \in I_P(\sigma)$ and $\psi_i \in I_P'(\sigma)$ the image of φ by the map in (i) for $i = 1, 2, 3$. Then ψ_1 is characterized by $\varphi(XT_{n_{Gw_P}}) = \psi_1(XT_{n_{Gw_P}})$ for any $X \in \mathcal{H}$. We denote the corresponding element to $\varphi \in I_P(\sigma)$ (resp. $\psi_i \in I_P'(\sigma)$) by $\varphi' \in (I_Q \circ I_P^Q)(\sigma)$ (resp. $\psi_i' \in (I_Q' \circ I_P^{Q'})(\sigma)$).

We consider ψ_2 . We have $\psi_2(XT_{n_{Gw_P}}) = \psi_2'(XT_{n_{Gw_P}})(1)$. Since $\ell(w_G w_P) = \ell(w_G w_Q) + \ell(w_Q w_P)$, we have $T_{n_{Gw_P}} = T_{n_{Gw_Q}} T_{n_{Qw_P}}$. Hence $\psi_2'(XT_{n_{Gw_P}})(1) = \psi_2'(XT_{n_{Gw_Q}} T_{n_{Qw_P}})(1)$. Since $T_{n_{Qw_P}}^Q \in \mathcal{H}_Q^-$ and $j_Q^-(T_{n_{Qw_P}}^Q) = T_{n_{Qw_P}}$ [Abe16, Lemma 2.6], we have $\psi_2'(XT_{n_{Gw_Q}} T_{n_{Qw_P}})(1) = \psi_2'(XT_{n_{Gw_Q}})(T_{n_{Qw_P}}^Q)$. Let

$\psi_2'' \in (I_Q' \circ I_P^Q)(\sigma)$ be the image of φ . Then $\psi_2''(XT_{n_{w_G w_Q}})(Y) = \varphi'(XT_{n_{w_G w_Q}})(Y)$ and $\psi_2''(X)(YT_{n_{w_Q w_P}}^Q) = \psi_2'(X)(YT_{n_{w_Q w_P}}^Q)$ for $X \in \mathcal{H}$ and $Y \in \mathcal{H}_Q$. Therefore we have $\psi_2(XT_{n_{w_G w_P}}) = \psi_2''(XT_{n_{w_G w_P}})(T_{n_{w_Q w_P}}^Q) = \varphi'(XT_{n_{w_G w_P}})(T_{n_{w_Q w_P}}^Q)$. Again, since $T_{n_{w_Q w_P}}^Q \in \mathcal{H}_Q^-$ and $j_Q^{-*}(T_{n_{w_Q w_P}}^Q) = T_{n_{w_Q w_P}}$ [Abe16, Lemma 2.6], we have $\varphi'(XT_{n_{w_G w_Q}})(T_{n_{w_Q w_P}}^Q) = \varphi'(XT_{n_{w_G w_Q}} T_{n_{w_Q w_P}})(1) = \varphi'(XT_{n_{w_G w_P}})(1) = \varphi(XT_{n_{w_G w_P}})$. Hence the map in (2) satisfies the characterization of the map in (1).

The proof for (3) is similar. Let $\psi_3'' \in (I_Q \circ I_P^{Q'})(\sigma)$ be the image of φ . Then we have

$$\begin{aligned} \psi_3(XT_{n_{w_G w_P}}) &= \psi_3'(XT_{n_{w_G w_P}})(1) \\ &= \psi_3'(XT_{n_{w_G w_Q}} T_{n_{w_Q w_P}})(1) \\ &= \psi_3'(XT_{n_{w_G w_Q}})(T_{n_{w_Q w_P}}^Q) \\ &= \psi_3''(XT_{n_{w_G w_Q}})(T_{n_{w_Q w_P}}^Q) \\ &= \varphi'(XT_{n_{w_G w_Q}})(T_{n_{w_Q w_P}}^Q) \\ &= \varphi'(XT_{n_{w_G w_Q}} T_{n_{w_Q w_P}})(1) \\ &= \varphi'(XT_{n_{w_G w_P}})(1) \\ &= \varphi(XT_{n_{w_G w_P}}). \end{aligned}$$

This finish the proof. $\square$

3.2. Steinberg modules. Next we consider the twist of the generalized Steinberg modules. Until subsection 3.7, we keep the following settings.

- P is a parabolic subgroup such that Δ_P is orthogonal to $\Delta \setminus \Delta_P$.
- Let P_2 be a parabolic subgroup corresponding to $\Delta \setminus \Delta_P$.
- σ is an $\mathcal{H}_P$ -module such that $P(\sigma) = G$.

We prove the following proposition.

Theorem 3.6. *Let Q be a parabolic subgroup containing P and Q^c the parabolic subgroup corresponding to $\Delta_P \cup (\Delta \setminus \Delta_Q)$. Then we have $\text{St}_Q(\sigma)^\iota \simeq \text{St}_{Q^c}(\sigma_{\ell-P}^{\iota_P})$.*

The proof of this proposition continues until subsection 3.7. In this subsection, we prove this proposition for $Q = P$.

Lemma 3.7. *Theorem 3.6 is true if $Q = P$.*

Proof. We use [Abe16, Proposition 3.12]. Let $w \in W_P(1)$ and assume that w is P -positive. Then $(\text{St}_P(\sigma))^\iota(T_w^*) = (-1)^{\ell(w)} \text{St}_P(\sigma)(T_w) = (-1)^{\ell(w)} \sigma(T_w^P) = (-1)^{\ell(w) - \ell_P(w)} \sigma^{\iota_P}(T_w^{P*}) = \sigma_{\ell-\ell_P}^{\iota_P}(T_w^{P*})$. On the other hand, for $w \in W_{\text{aff}, P_2}(1)$, we have $(\text{St}_P(\sigma))^\iota(T_w^*) = (-1)^{\ell(w)} \text{St}_P(\sigma)(T_w) = 1$. Therefore, by a characterization of the extension, we have $(\text{St}_P(\sigma))^\iota \simeq e_G(\sigma_{\ell-\ell_P}^{\iota_P}) = \text{St}_G(\sigma_{\ell-\ell_P}^{\iota_P})$. Since $P^c = G$, we get the lemma. $\square$

3.3. An exact sequence. We express $\text{St}_{Q^c}(\sigma)$ as the kernel of a certain homomorphism. As a consequence, we deduce Theorem 3.6 to the exactness of a certain sequence (Lemma 3.9).

Let Q be a parabolic subgroup containing P . Then we have an exact sequence

$$0 \rightarrow \sum_{Q \supsetneq R \supsetneq P} I_R^Q(e_R(\sigma)) \rightarrow I_P^Q(\sigma) \rightarrow \text{St}_P^Q(\sigma) \rightarrow 0.$$

Applying I_Q and using the transitivity [Vig15b, Proposition 4.10], we have an exact sequence

$$0 \rightarrow \sum_{Q \supset R \supsetneq P} I_R(e_R(\sigma)) \rightarrow I_P(\sigma) \rightarrow I_Q(\text{St}_P^Q(\sigma)) \rightarrow 0.$$

Let Q_1 be a parabolic subgroup containing Q . Then we have $\sum_{Q \supset R \supsetneq P} I_R(e_R(\sigma)) \subset \sum_{Q_1 \supset R \supsetneq P} I_R(e_R(\sigma))$. Hence we have a homomorphism $I_Q(\text{St}_P^Q(\sigma)) \rightarrow I_{Q_1}(\text{St}_P^{Q_1}(\sigma))$ which makes the following diagram commutative.

$$\begin{array}{ccccccc} 0 & \longrightarrow & \sum_{Q \supset R \supsetneq P} I_R(e_R(\sigma)) & \longrightarrow & I_P(\sigma) & \longrightarrow & I_Q(\text{St}_P^Q(\sigma)) \longrightarrow 0 \\ & & \downarrow & & \parallel & & \downarrow \\ 0 & \longrightarrow & \sum_{Q_1 \supset R \supsetneq P} I_R(e_R(\sigma)) & \longrightarrow & I_P(\sigma) & \longrightarrow & I_{Q_1}(\text{St}_P^{Q_1}(\sigma)) \longrightarrow 0. \end{array}$$

Lemma 3.8. *With the above homomorphisms, we have an exact sequence*

$$0 \rightarrow \text{St}_{Q^c}(\sigma) \rightarrow I_Q(\text{St}_P^Q(\sigma)) \rightarrow \bigoplus_{Q_1 \supsetneq P} I_{Q_1}(\text{St}_P^{Q_1}(\sigma)).$$

We will prove this lemma at the end of this subsection. Applying ι to the exact sequence in the lemma, we have

$$0 \rightarrow \text{St}_{Q^c}(\sigma)^\iota \rightarrow I_Q(\text{St}_P^Q(\sigma))^\iota \rightarrow \bigoplus_{Q_1 \supsetneq P} I_{Q_1}(\text{St}_P^{Q_1}(\sigma))^\iota.$$

Using Lemma 3.7 and [Abe16, Lemma 4.9, Proposition 4.11], we have $I_Q(\text{St}_P^Q(\sigma))^\iota = I'_Q((\text{St}_P^Q(\sigma))^{\iota_Q}) = I'_Q(e_Q(\sigma_{\ell_Q - \ell_P}^{\iota_P})) = I'_Q(e_Q(\sigma_{\ell - \ell_P}^{\iota_P}))$. We get the following exact sequence

$$0 \rightarrow (\text{St}_{Q^c}(\sigma))^\iota \rightarrow I'_Q(e_Q(\sigma_{\ell - \ell_P}^{\iota_P})) \rightarrow \bigoplus_{Q_1 \supsetneq Q} I'_{Q_1}(e_{Q_1}(\sigma_{\ell - \ell_P}^{\iota_P})).$$

Therefore Theorem 3.6 follows from the following lemma.

Lemma 3.9. *The following sequence is exact.*

$$\bigoplus_{Q_1 \supsetneq Q} I_{Q_1}(e_{Q_1}(\sigma)) \rightarrow I_Q(e_Q(\sigma)) \rightarrow I'_Q(e_Q(\sigma)) \rightarrow \bigoplus_{Q_1 \supsetneq Q} I'_{Q_1}(e_{Q_1}(\sigma)).$$

Here $I_Q(e_Q(\sigma)) \rightarrow I'_Q(e_Q(\sigma))$ is given in Proposition 3.1 and $I_{Q_1}(e_{Q_1}(\sigma)) \rightarrow I'_Q(e_Q(\sigma))$ is the natural embedding.

As the end of this subsection, we prove Lemma 3.8. We need the following lemma. For parabolic subgroups $P_1, \dots, P_r$, let $\langle P_1, \dots, P_r \rangle$ be the parabolic subgroup generated by $P_1, \dots, P_r$. In other words, $\langle P_1, \dots, P_r \rangle$ is the parabolic subgroup corresponding to $\Delta_{P_1} \cup \dots \cup \Delta_{P_r}$.

Lemma 3.10. *Let $\mathcal{P}_1, \mathcal{P}_2 \subset \{Q \mid Q \supset P\}$. Then, in $I_P(\sigma)$, we have*

$$\left(\sum_{Q_1 \in \mathcal{P}_1} I_{Q_1}(e_{Q_1}(\sigma)) \right) \cap \left(\sum_{Q_2 \in \mathcal{P}_2} I_{Q_2}(e_{Q_2}(\sigma)) \right) = \sum_{Q_1 \in \mathcal{P}_1, Q_2 \in \mathcal{P}_2} I_{\langle Q_1, Q_2 \rangle}(e_{\langle Q_1, Q_2 \rangle}(\sigma)).$$

Remark 3.11. We prove Lemma 3.8 and 3.10 for any commutative ring C .

Proof. Note that Δ_P is orthogonal to Δ_{P_2} as we assumed. Therefore we have $W_0^P = W_{0,P_2}$ and for any parabolic subgroup Q containing P , we have $W_{0,P_2} = W_0^Q W_{0,Q \cap P_2}$.

Put $I_Q = I_Q(e_Q(\sigma))$. We prove the lemma by induction on $\#\mathcal{P}_1$. Assume that $\#\mathcal{P}_1 = 1$. By [Abe16, Lemma 3.8], it is sufficient to prove that $I_{Q_1} \cap I_{Q_2} = I_{\langle Q_1, Q_2 \rangle}$. Obviously we have $I_{Q_1} \cap I_{Q_2} \supset I_{\langle Q_1, Q_2 \rangle}$. Let $\varphi \in I_{Q_1} \cap I_{Q_2}$. Then for $w \in W_0^{\langle Q_1, Q_2 \rangle}$ and $\alpha \in \Delta_{Q_1} \setminus \Delta_P$, we have $\varphi(T_{n_{ws_\alpha}}) = \varphi(T_{n_w})e_{Q_1}(\sigma)(T_{n_{s_\alpha}}^{Q_1})$. Since $n_{s_\alpha} \in W_{P_2 \cap Q, \text{aff}}$, we have $e_{Q_1}(\sigma)(T_{n_{s_\alpha}}^{Q_1}) = q_{s_\alpha}$. Therefore we have $\varphi(T_{n_{ws_\alpha}}) = q_{s_\alpha} \varphi(T_{n_w})$. This also holds for $\alpha \in \Delta_{Q_2} \setminus \Delta_P$. Therefore $\varphi(T_{n_{wv}}) = q_v \varphi(T_{n_w})$ for any v generated by $\{s_\alpha \mid \alpha \in (\Delta_{Q_1} \cup \Delta_{Q_2}) \setminus \Delta_P\}$.

Since $(\Delta_{Q_1} \cup \Delta_{Q_2}) \setminus \Delta_P = \Delta_{\langle Q_1, Q_2 \rangle \cap P_2}$, the group generated by $\{s_\alpha \mid \alpha \in (\Delta_{Q_1} \cup \Delta_{Q_2}) \setminus \Delta_P\}$ is $W_{0, \langle Q_1, Q_2 \rangle \cap P_2}$. Hence $\varphi(T_{n_{wv}}) = q_v \varphi(T_{n_w})$ for any $w \in W_0^{\langle Q_1, Q_2 \rangle}$ and $v \in W_{0, \langle Q_1, Q_2 \rangle \cap P_2}$. Define $\varphi' \in I_{\langle Q_1, Q_2 \rangle}(e_{\langle Q_1, Q_2 \rangle}(\sigma))$ by $\varphi'(T_{n_w}) = \varphi(T_{n_w})$ for any $w \in W_0^{\langle Q_1, Q_2 \rangle}$. (Such element uniquely exists by Proposition 2.3.) Then φ' also satisfies $\varphi(T_{n_{wv}}) = q_v \varphi(T_{n_w})$ for any $w \in W_0^{\langle Q_1, Q_2 \rangle}$ and $v \in W_{0, \langle Q_1, Q_2 \rangle \cap P_2}$. Hence $\varphi(T_{n_w}) = \varphi'(T_{n_w})$ for any $w \in W_0^{\langle Q_1, Q_2 \rangle} W_{0, \langle Q_1, Q_2 \rangle \cap P_2} = W_{0,P_2} = W_0^P$. By Proposition 2.3, $\varphi = \varphi' \in I_{\langle Q_1, Q_2 \rangle}$.

Now we prove the general case. Obviously we have

$$\left(\sum_{Q_1 \in \mathcal{P}_1} I_{Q_1} \right) \cap \left(\sum_{Q_2 \in \mathcal{P}_2} I_{Q_2} \right) \supset \sum_{Q_1 \in \mathcal{P}_1, Q_2 \in \mathcal{P}_2} I_{\langle Q_1, Q_2 \rangle}.$$

We prove the reverse inclusion. Take f from the left hand side. Fix $Q_0 \in \mathcal{P}_1$ and put $\mathcal{P}'_1 = \mathcal{P}_1 \setminus \{Q_0\}$. Take $f_1 \in I_{Q_0}$ and $f_2 \in \sum_{Q_1 \in \mathcal{P}'_1} I_{Q_1}$ such that $f = f_1 + f_2$. Then we have

$$f_2 \in \left(\sum_{Q_2 \in \mathcal{P}_2} I_{Q_2} + I_{Q_0} \right) \cap \sum_{Q_1 \in \mathcal{P}'_1} I_{Q_1}.$$

By inductive hypothesis, the right hand side is

$$\sum_{Q_1 \in \mathcal{P}'_1, Q_2 \in \mathcal{P}_2} I_{\langle Q_1, Q_2 \rangle} + \sum_{Q_1 \in \mathcal{P}'_1} I_{\langle Q_0, Q_1 \rangle}.$$

Since $I_{\langle Q_0, Q_1 \rangle} \subset I_{Q_0}$, we get

$$f_2 \in \sum_{Q_1 \in \mathcal{P}'_1, Q_2 \in \mathcal{P}_2} I_{\langle Q_1, Q_2 \rangle} + I_{Q_0}.$$

We have $f_1 \in I_{Q_0}$. Therefore

$$f = f_1 + f_2 \in \sum_{Q_1 \in \mathcal{P}'_1, Q_2 \in \mathcal{P}_2} I_{\langle Q_1, Q_2 \rangle} + I_{Q_0}.$$

Take $f'_1 \in \sum_{Q_1 \in \mathcal{P}'_1, Q_2 \in \mathcal{P}_2} I_{\langle Q_1, Q_2 \rangle}$ and $f'_2 \in I_{Q_0}$ such that $f = f'_1 + f'_2$. Then $f'_1 \in \sum_{Q_2 \in \mathcal{P}_2} I_{Q_2}$. By the assumption, $f \in \sum_{Q_2 \in \mathcal{P}_2} I_{Q_2}$. Therefore we have $f'_2 = f - f'_1 \in \sum_{Q_2 \in \mathcal{P}_2} I_{Q_2}$. Hence

$$f'_2 \in I_{Q_0} \cap \sum_{Q_2 \in \mathcal{P}_2} I_{Q_2} = \sum_{Q_2 \in \mathcal{P}_2} I_{\langle Q_0, Q_2 \rangle}.$$

Here we use the lemma for $\mathcal{P}_1 = \{Q_0\}$. Hence

$$\begin{aligned} f &= f'_1 + f'_2 \in \sum_{Q_1 \in \mathcal{P}'_1, Q_2 \in \mathcal{P}_2} I_{\langle Q_1, Q_2 \rangle} + \sum_{Q_2 \in \mathcal{P}_2} I_{\langle Q_0, Q_2 \rangle} \\ &= \sum_{Q_1 \in \mathcal{P}_1, Q_2 \in \mathcal{P}_2} I_{\langle Q_1, Q_2 \rangle}. \end{aligned}$$

We get the lemma. $\square$

Proof of Lemma 3.8. Let $Q_1 \supsetneq Q$. By the exact sequence before Lemma 3.8, the kernel of $I_P(\sigma) \rightarrow I_Q(\text{St}_P^Q(\sigma)) \rightarrow I_{Q_1}(\text{St}_P^{Q_1}(\sigma))$ is $\sum_{Q_1 \supset R \supsetneq P} I_R(e_R(\sigma))$. Since $\text{Ker}(I_P(\sigma) \rightarrow I_Q(\text{St}_P^Q(\sigma))) = \sum_{Q \supset R \supsetneq P} I_R(e_R(\sigma))$, we have

$$\begin{aligned} &\text{Ker} \left(I_Q(\text{St}_P^Q(\sigma)) \rightarrow \bigoplus_{Q_1 \supsetneq Q} I_{Q_1}(\text{St}_P^{Q_1}(\sigma)) \right) \\ &\simeq \text{Ker} \left(I_P(\sigma) \rightarrow \bigoplus_{Q_1 \supsetneq Q} I_{Q_1}(\text{St}_P^{Q_1}(\sigma)) \right) \Big/ \text{Ker}(I_P(\sigma) \rightarrow I_Q(\text{St}_P^Q(\sigma))) \\ &= \left(\bigcap_{Q_1 \supsetneq Q} \text{Ker}(I_P(\sigma) \rightarrow I_{Q_1}(\text{St}_P^{Q_1}(\sigma))) \right) \Big/ \text{Ker}(I_P(\sigma) \rightarrow I_Q(\text{St}_P^Q(\sigma))) \\ &= \bigcap_{Q_1 \supsetneq Q} \sum_{Q_1 \supset R \supsetneq P} I_R(e_R(\sigma)) \Big/ \sum_{Q \supset R \supsetneq P} I_R(e_R(\sigma)). \end{aligned}$$

Set $A = \bigcap_{Q_1 \supsetneq Q} \sum_{Q_1 \supset R \supsetneq P} I_R(e_R(\sigma))$ and $B = \sum_{Q \supset R \supsetneq P} I_R(e_R(\sigma))$. We prove the following.

- (1) $I_{Q^c}(e_{Q^c}(\sigma)) + B = A$.
- (2) $I_{Q^c}(e_{Q^c}(\sigma)) \cap B = \sum_{R \supsetneq Q^c} I_R(e_R(\sigma))$.

First we prove (1). We prove $I_{Q^c}(e_{Q^c}(\sigma)) + B \subset A$. Let $Q_1 \supsetneq Q$. Take $\alpha \in \Delta_{Q_1} \setminus \Delta_Q$ and let R be a parabolic subgroup corresponding to $\Delta_P \cup \{\alpha\}$. Since $\alpha \in \Delta_{Q^c}$ and $\Delta_P \subset \Delta_{Q^c}$, we have $\Delta_R \subset \Delta_{Q^c}$. Hence $I_{Q^c}(e_{Q^c}(\sigma)) \subset I_R(e_R(\sigma)) \subset A$. Obviously we have $B \subset A$.

We prove the reverse inclusion. By Lemma 3.10, we have

$$A = \sum_{(R_{Q_1})_{Q_1 \supsetneq Q}} I_{\langle R_{Q_1} \rangle_{Q_1}}(e_{\langle R_{Q_1} \rangle_{Q_1}}(\sigma)),$$

where R_{Q_1} satisfies $Q_1 \supset R_{Q_1} \supsetneq P$ and $\langle R_{Q_1} \rangle_{Q_1}$ is the group generated by $\{R_{Q_1} \mid Q_1 \supsetneq Q\}$. Hence it is sufficient to prove that each $I_{\langle R_{Q_1} \rangle_{Q_1}}(e_{\langle R_{Q_1} \rangle_{Q_1}}(\sigma))$ is contained in $I_{Q^c}(e_{Q^c}(\sigma)) + B$. If $Q \supset R_{Q_0}$ for some $Q_0 \supsetneq Q$, then for such Q_0 , we have $I_{\langle R_{Q_1} \rangle_{Q_1}}(e_{\langle R_{Q_1} \rangle_{Q_1}}(\sigma)) \subset I_{R_{Q_0}}(e_{R_{Q_0}}(\sigma)) \subset B$. Assume that $Q \not\supset R_{Q_1}$ for any $Q_1 \supsetneq Q$ and we prove $\Delta_{Q^c} \subset \bigcup_{Q_1 \supsetneq Q} \Delta_{R_{Q_1}}$. Let $\alpha \in \Delta_{Q^c} = (\Delta \setminus \Delta_Q) \cup \Delta_P$. If $\alpha \in \Delta_P$, then we have $\alpha \in \bigcup_{Q_1 \supsetneq Q} \Delta_{R_{Q_1}}$ since $P \subset R_{Q_1}$. Assume that $\alpha \in \Delta \setminus \Delta_Q$. Let Q_α be the parabolic subgroup corresponding to $\Delta_Q \cup \{\alpha\}$. Then by the assumption, $\Delta_{R_{Q_\alpha}}$ is contained in $\Delta_{Q_\alpha} = \Delta_Q \cup \{\alpha\}$ and is not contained in Δ_Q . Hence $\alpha \in \Delta_{R_{Q_\alpha}}$. Therefore $\alpha \in \Delta_{R_{Q_\alpha}} \subset \bigcup_{Q_1 \supsetneq Q} \Delta_{R_{Q_1}} = \Delta_{\langle R_{Q_1} \rangle_{Q_1}}$ by taking $Q_1 = Q_\alpha$. Hence we have $Q^c \subset \langle R_{Q_1} \rangle_{Q_1}$. Therefore $I_{\langle R_{Q_1} \rangle_{Q_1}}(e_{\langle R_{Q_1} \rangle_{Q_1}}(\sigma)) \subset I_{Q^c}(e_{Q^c}(\sigma))$. We get (1).

We prove (2). By the above lemma again, we have

$$I_{Q^c}(e_{Q^c}(\sigma)) \cap B = \sum_{Q \supset R \supseteq P} I_{\langle R, Q^c \rangle}(e_{\langle R, Q^c \rangle}(\sigma)).$$

First we prove $I_{Q^c}(e_{Q^c}(\sigma)) \cap B \subset \sum_{R_1 \supseteq Q^c} I_{R_1}(e_{R_1}(\sigma))$, namely, for each R such that $Q \supset R \supseteq P$ we have $I_{\langle R, Q^c \rangle}(e_{\langle R, Q^c \rangle}(\sigma)) \subset \sum_{R_1 \supseteq Q^c} I_{R_1} R(e_{R_1}(\sigma))$. For such R , we can take $\alpha \in \Delta_R \setminus \Delta_P$. Since $\alpha \in \Delta_Q \setminus \Delta_P$, $\alpha \notin \Delta_{Q^c}$. Hence $\Delta_{\langle R, Q^c \rangle} = \Delta_R \cup \Delta_{Q^c} \supset \{\alpha\} \cup \Delta_{Q^c} \supseteq \Delta_{Q^c}$. Therefore $\langle R, Q^c \rangle \supseteq Q^c$. Hence $I_{\langle R, Q^c \rangle}(e_{\langle R, Q^c \rangle}(\sigma)) \subset \sum_{R_1 \supseteq Q^c} I_{R_1}(e_{R_1}(\sigma))$ by taking $R_1 = \langle R, Q^c \rangle$.

We prove $I_R(e_R(\sigma)) \subset I_{Q^c}(e_{Q^c}(\sigma)) \cap B$ for any R such that $R \supseteq Q^c$. We can take $\alpha \in \Delta_R \setminus \Delta_{Q^c}$. Let P_α be the parabolic subgroup corresponding to $\Delta_P \cup \{\alpha\}$. Since $\alpha \notin \Delta_{Q^c}$, we have $\alpha \in \Delta_Q$. Therefore $Q \supset P_\alpha \supseteq P$. Hence $\Delta_R \supset \Delta_{Q^c} \cup \{\alpha\} = \Delta_{\langle P_\alpha, Q^c \rangle}$. Therefore $R \supset \langle P_\alpha, Q^c \rangle$. Hence $I_R(e_R(\sigma)) \subset I_{\langle P_\alpha, Q^c \rangle}(e_{\langle P_\alpha, Q^c \rangle}(\sigma)) \subset I_{Q^c}(e_{Q^c}(\sigma)) \cap B$. We get (2) and the proof of the lemma is finished. $\square$

3.4. The kernel of $I_Q(e_Q(\sigma)) \rightarrow I'_Q(e_Q(\sigma))$. We determine $\text{Ker}(I_Q(e_Q(\sigma)) \rightarrow I'_Q(e_Q(\sigma)))$, namely we prove the following lemma.

Lemma 3.12. *Set $A = \{w \in W_0^Q \mid \Delta_w = \Delta_Q\}$. Then we have*

$$\begin{aligned} & \text{Ker}(I_Q(e_Q(\sigma)) \rightarrow I'_Q(e_Q(\sigma))) \\ &= \{\varphi \in I_Q(e_Q(\sigma)) \mid \varphi(XT_{n_w}) = 0 \text{ for any } X \in \mathcal{H} \text{ and } w \in A\} \\ &= \{\varphi \in I_Q(e_Q(\sigma)) \mid \varphi(T_{n_w}) = 0 \text{ for any } w \in A\} = \sum_{Q_1 \supseteq Q} I_{Q_1}(e_{Q_1}(\sigma)). \end{aligned}$$

The last equality follows from Lemma 2.6 and the following lemma.

Lemma 3.13. *We have $\bigcap_{Q_1 \supseteq Q} (W_0^Q \setminus W_0^{Q_1}) = \{w \in W_0^Q \mid \Delta_w = \Delta_Q\}$.*

Proof. Let $w \in W_0^Q$. Then $w(\Delta_Q) > 0$ and we have $\Delta_w = \Delta_Q$ if and only if for any $Q_1 \supseteq Q$, $w(\Delta_{Q_1}) \not\subset \Sigma^+$. Since we have $w(\Delta_{Q_1}) \not\subset \Sigma^+$ if and only if $w \notin W_0^{Q_1}$, we get the lemma. $\square$

Let $\varphi \in I_Q(e_Q(\sigma))$ such that $\varphi(T_{n_w}) = 0$ for any $w \in A$ and take $X \in \mathcal{H}$. The last equality implies that the set of such φ is stable under $\mathcal{H}$. Hence φX also satisfies the same condition. Therefore $\varphi(XT_{n_w}) = 0$ for any $w \in A$. Namely we get the second equality.

Let ψ be the image of φ under $I_Q(e_Q(\sigma)) \rightarrow I'_Q(e_Q(\sigma))$. Then ψ is characterized by $\varphi(XT_{n_{w_G w_Q}}) = \psi(XT_{n_{w_G w_Q}})$. Therefore we have

$$\begin{aligned} & \text{Ker}(I_Q(e_Q(\sigma)) \rightarrow I'_Q(e_Q(\sigma))) \\ &= \{\varphi \in I_Q(e_Q(\sigma)) \mid \varphi(XT_{n_{w_G w_Q}}) = 0 \text{ for any } X \in \mathcal{H}\}. \end{aligned}$$

Therefore, to prove the lemma, it is sufficient to prove the following lemma.

Lemma 3.14. *Let $\varphi \in I_Q(e_Q(\sigma))$. Assume that $\varphi(XT_{n_{w_G w_Q}}) = 0$ for any $X \in \mathcal{H}$. Then we have $\varphi(XT_{n_w}) = 0$ for any $X \in \mathcal{H}$ and $w \in W_0^Q$ such that $\Delta_w = \Delta_Q$.*

Proof. We prove the lemma by backward induction on $\ell(w)$. If $w \neq w_G w_Q$, then there exists $\alpha \in \Delta$ such that $s_\alpha w > w$, $\Delta_w = \Delta_{s_\alpha w}$ and $w^{-1}(\alpha)$ is not simple [Abe, Lemma 3.15]. Set $s = s_\alpha$. Since $\Delta_{sw} = \Delta_w = \Delta_Q$, we have $sw \in W_0^Q$. If

$w^{-1}(\alpha) \in \Sigma_Q^+$, then since $sw \in W_0^Q$, we have $-\alpha = sw(w^{-1}(\alpha)) \in \Sigma^+$. This is a contradiction. Hence

$$w^{-1}(\alpha) \in \Sigma^+ \setminus \Sigma_Q^+.$$

Take $\lambda_P^- \in Z(W_P(1))$ as in Proposition 2.2. Put $\lambda = n_w \cdot (\lambda_P^-)^2$. We prove:

Claim. $E_-(\lambda n_s^{-1})(T_{n_s} - c_{n_s}) = E_{o_-}(\lambda)$ in $\mathcal{H}$

We calculate the left hand side in $\mathcal{H}[q_s^{\pm 1}]$. We use notation in [Abe, Lemma 2.10]. Since $w^{-1}(\alpha) \in \Sigma^+ \setminus \Sigma_Q^+ \subset \Sigma^+ \setminus \Sigma_P^+$, we have $\langle \alpha, \nu(\lambda) \rangle = \langle w^{-1}(\alpha), \nu((\lambda_P^-)^2) \rangle > 0$. Therefore we have $\ell(\lambda n_s^{-1}) = \ell(\lambda) - 1$ by [Abe16, Lemma 2.17]. Hence $q_{\lambda n_s^{-1}} = q_\lambda q_{n_s}^{-1}$. Therefore we have

$$\begin{aligned} E_-(\lambda n_s^{-1}) &= E_-(n_s^{-1}(n_s \cdot \lambda)) \\ &= q_{\lambda n_s^{-1}}^{1/2} q_{n_s}^{-1/2} T_{n_s^{-1}}^* \theta(n_s \cdot \lambda) \\ &= q_\lambda^{1/2} q_{n_s}^{-1} T_{n_s^{-1}}^* \theta(n_s \cdot \lambda). \end{aligned}$$

Hence by [Abe, Lemma 2.10], we have

$$\begin{aligned} (3.1) \quad & E_-(\lambda n_s^{-1}) T_{n_s} \\ &= q_\lambda^{1/2} q_{n_s}^{-1} T_{n_s^{-1}}^* \theta(n_s \cdot \lambda) T_{n_s} \\ &= q_\lambda^{1/2} q_{n_s}^{-1} T_{n_s^{-1}}^* T_{n_s} \theta(\lambda) + \sum_{k=0}^{\langle \alpha, \nu(\lambda) \rangle - 1} q_\lambda^{1/2} q_{n_s}^{-1} T_{n_s^{-1}}^* \theta(n_s \cdot \lambda \mu_{n_s}(k)) c_{n_s, k}. \end{aligned}$$

We have

$$q_\lambda^{1/2} q_{n_s}^{-1} T_{n_s^{-1}}^* T_{n_s} \theta(\lambda) = q_\lambda^{1/2} \theta(\lambda) = E_{o_-}(\lambda).$$

and if $k = 0$, since $q_{\lambda n_s^{-1}} = q_\lambda q_{n_s}^{-1}$, we have

$$\begin{aligned} q_\lambda^{1/2} q_{n_s}^{-1} T_{n_s^{-1}}^* \theta(n_s \cdot \lambda \mu_{n_s}(k)) c_{n_s, k} &= q_\lambda^{1/2} q_{n_s}^{-1} T_{n_s^{-1}}^* \theta(n_s \cdot \lambda) c_{n_s} \\ &= q_{\lambda n_s^{-1}}^{1/2} q_{n_s}^{-1/2} T_{n_s^{-1}}^* \theta(n_s \cdot \lambda) c_{n_s} \\ &= E_-(n_s^{-1}(n_s \cdot \lambda)) c_{n_s} = E_-(\lambda n_s^{-1}) c_{n_s}. \end{aligned}$$

We prove that if $1 \leq k \leq \langle \alpha, \nu(\lambda) \rangle - 1$, then $q_\lambda^{1/2} q_{n_s}^{-1} T_{n_s^{-1}}^* \theta(n_s \cdot \lambda \mu_{n_s}(k)) c_{n_s, k} = 0$ in $\mathcal{H}$. We have

$$\begin{aligned} & q_\lambda^{1/2} q_{n_s}^{-1} T_{n_s^{-1}}^* \theta(n_s \cdot \lambda \mu_{n_s}(k)) c_{n_s, k} \\ &= q_\lambda^{1/2} q_{n_s^{-1}(n_s \cdot \lambda \mu_{n_s}(k))}^{-1/2} q_{n_s}^{-1/2} E_-(n_s^{-1}(n_s \cdot \lambda \mu_{n_s}(k))) c_{n_s, k}. \end{aligned}$$

Hence (3.1) is an expansion of $E_-(\lambda n_s^{-1}) T_{n_s}$ with respect to the basis $\{E_-(w) \mid w \in W(1)\}$. Since this is a basis of $\mathcal{H}[q_s]$ as a $C[q_s]$ -module, each coefficient is in $C[q_s]$. Hence $q_\lambda^{1/2} q_{n_s^{-1}(n_s \cdot \lambda \mu_{n_s}(k))}^{-1/2} q_{n_s}^{-1/2} \in C[q_s]$. Namely, for each $s \in S_{\text{aff}}$ there exists $k_s \in \mathbb{Z}_{\geq 0}/\sim$ (where the equivalence relation $\sim$ is defined by the adjoint action of W on S_{aff}) such that $q_\lambda^{1/2} q_{n_s^{-1}(n_s \cdot \lambda \mu_{n_s}(k))}^{-1/2} q_{n_s}^{-1/2} = \prod_{s \in S_{\text{aff}}/\sim} q_s^{k_s}$. We have $\sum_s k_s = (1/2)(\ell(\lambda) - \ell(n_s^{-1}(n_s \cdot \lambda \mu_{n_s}(k))) - \ell(n_s))$. We calculate

$$\begin{aligned} \ell(\lambda) - \ell(n_s^{-1}(n_s \cdot \lambda \mu_{n_s}(k))) - \ell(n_s) &\geq \ell(\lambda) - \ell(n_s^{-1}) - \ell(n_s \cdot \lambda \mu_{n_s}(k)) - \ell(n_s) \\ &= \ell(\lambda) - \ell(n_s \cdot \lambda \mu_{n_s}(k)) - 2. \end{aligned}$$

By [Abe, Lemma 2.12], $\ell(\lambda) - \ell(n_s \cdot \lambda \mu_{n_s}(k)) \geq 2 \min\{k, \langle \alpha, \nu(\lambda) \rangle - k\}$. If the equality holds, again by [Abe, Lemma 2.12], there exists $v \in W_0$ such that $v\nu(\lambda)$ is dominant and $v(\alpha)$ is simple.

Assume that $v\nu(\lambda)$ is dominant for $v \in W_0$. We have $\nu(\lambda) = w(\nu((\lambda_P^-)^2))$ and since $\nu((\lambda_P^-)^2)$ is dominant, we have $vw \in \text{Stab}_{W_0}(\nu((\lambda_P^-)^2))$. By the condition of λ_P^- , the stabilizer of $\nu((\lambda_P^-)^2)$ is $W_{0,P}$. Hence $vw \in W_{0,P}$. Since $w^{-1}(\alpha) \in \Sigma^+ \setminus \Sigma_Q^+$, we have $w^{-1}(\alpha) \in \Sigma_{P_2}^+$. Any element in Σ_{P_2} is fixed by elements in $W_{0,P}$. Hence $vw(w^{-1}(\alpha)) = w^{-1}(\alpha)$. Therefore $v(\alpha) = w^{-1}(\alpha)$. This is not simple by the condition on α .

Hence we always have

$$\ell(\lambda) - \ell(n_s^{-1}(n_s \cdot \lambda \mu_{n_s}(k))) - \ell(n_s) > 0$$

for $1 \leq k \leq \langle \alpha, \nu(\lambda) \rangle - 1$. Hence $\sum_s k_s > 0$. Therefore there exists s such that $k_s > 0$. Hence $\prod_s q_s^{k_s} = 0$ in $\mathcal{H}$. We get $q_\lambda^{1/2} q_{n_s}^{-1} T_{n_s^{-1}}^* \theta(n_s \cdot \lambda \mu_{n_s}(k)) c_{n_s, k} = 0$. Therefore we have

$$E_-(\lambda n_s^{-1}) T_{n_s} = E_{o_-}(\lambda) + E_-(\lambda n_s^{-1}) c_{n_s}.$$

This gives the claim.

We return to the proof of the lemma. Since $n_w^{-1} \cdot \lambda = (\lambda_P^-)^2$ is P -negative, by Lemma 2.7, we have

$$\varphi(X E_{o_-}(\lambda) T_{n_w}) = \varphi(X T_{n_w}) \sigma(E_{o_-, P}^P(n_w^{-1} \cdot \lambda)).$$

Since $\sigma(E_{o_-, P}^P(n_w^{-1} \cdot \lambda)) = \sigma(E_{o_-, P}^P((\lambda_0^-)^2))$ is invertible, it is sufficient to prove that $\varphi(X E_{o_-}(\lambda) T_{n_w}) = 0$. By the claim, we have

$$\varphi(X E_{o_-}(\lambda) T_{n_w}) = \varphi(X E_-(\lambda n_s^{-1}) T_{n_s} T_{n_w}) - \varphi(X E_-(\lambda n_s^{-1}) c_{n_s} T_{n_w})$$

By inductive hypothesis, $\varphi(X E_-(\lambda n_s^{-1}) T_{n_s} T_{n_w}) = \varphi(X E_-(\lambda n_s^{-1}) T_{n_{sw}}) = 0$. We have

$$\varphi(X E_{o_-}(\lambda n_s^{-1}) c_{n_s} T_{n_w}) = \varphi(X((\lambda n_s^{-1}) \cdot c_{n_s}) E_-(\lambda n_s^{-1}) T_{n_w})$$

Set $\lambda' = n_w \cdot \lambda_P^-$. We have $\ell(\lambda n_s^{-1}) = \ell(\lambda) - 1 = \ell((\lambda')^2) - 1 = 2\ell(\lambda') - 1$ as $\langle \alpha, \nu(\lambda) \rangle > 0$. Since $\langle \alpha, \nu(\lambda') \rangle = \langle w^{-1}(\alpha), \nu(\lambda_P^-) \rangle > 0$, we have $\ell(\lambda') - 1 = \ell(\lambda' n_s^{-1})$. By [Abe16, Lemma 2.15], we have $\ell(\lambda') = \ell(n_s \cdot \lambda')$. Hence $\ell(\lambda n_s^{-1}) = \ell(\lambda' n_s^{-1}) + \ell(n_s \cdot \lambda')$. Therefore

$$E_-(\lambda n_s^{-1}) = E_-(\lambda' n_s^{-1} (n_s \cdot \lambda')) = E_-(\lambda' n_s^{-1}) E_{o_-}(n_s \cdot \lambda')$$

by the definition of $E_-(\lambda n_s^{-1})$. Hence

$$\begin{aligned} & \varphi(X((\lambda n_s^{-1}) \cdot c_{n_s}) E_-(\lambda n_s^{-1}) T_{n_w}) \\ &= \varphi(X((\lambda n_s^{-1}) \cdot c_{n_s}) E_-(\lambda' n_s^{-1}) E_{o_-}(n_s \cdot \lambda') T_{n_w}). \end{aligned}$$

Since $w^{-1}(\alpha) \in \Sigma^+ \setminus \Sigma_Q^+ \subset \Sigma^+ \setminus \Sigma_P^+$, we have $\langle w^{-1}(\alpha), \nu(n_w^{-1} n_s \cdot \lambda') \rangle = -\langle \alpha, \nu(\lambda') \rangle = -\langle w^{-1}(\alpha), \nu(\lambda_P^-) \rangle < 0$. Therefore $n_w^{-1} n_s \cdot \lambda'$ is not Q -negative. Hence $\varphi(X((\lambda n_s^{-1}) \cdot c_{n_s}) E_-(\lambda' n_s^{-1}) E_{o_-}(n_s \cdot \lambda') T_{n_w}) = 0$ by Proposition 2.4. $\square$

3.5. The homomorphism $I'_Q(e_Q(\sigma)) \rightarrow I'_{Q_1}(e_{Q_1}(\sigma))$. Let $Q_1 \supset Q \supset P$ be parabolic subgroups. Recall that we have the homomorphism $I'_Q(e_Q(\sigma)) \rightarrow I'_{Q_1}(e_{Q_1}(\sigma))$. This is defined by $I_Q(\text{St}_P^Q(\sigma_{\ell-P}^{\iota_P})) \rightarrow I_{Q_1}(\text{St}_P^{Q_1}(\sigma_{\ell-P}^{\iota_P}))$ with ι . We give the following description of this homomorphism.

Proposition 3.15. *Let $\varphi \in I'_Q(e_Q(\sigma))$ and $\varphi' \in I'_{Q_1}(e_{Q_1}(\sigma))$ be the image of φ . Then for $w \in W_0^{Q_1}$, we have $\varphi'(T_{n_w}^*) = (-1)^{\ell(w_{Q_1}w_Q)}\varphi(T_{n_{w_{Q_1}w_Q}}^*)$. In particular, combining with [Abe16, Proposition 4.12], we have*

$$\begin{aligned} & \text{Ker}(I'_Q(e_Q(\sigma)) \rightarrow I'_{Q_1}(e_{Q_1}(\sigma))) \\ &= \{\varphi \in I'_Q(e_Q(\sigma)) \mid \varphi(T_{n_w}^*) = 0 \text{ for any } w \in W_0^{Q_1}w_{Q_1}w_Q\}. \end{aligned}$$

First we describe the homomorphism $I_Q(\text{St}_P^Q(\sigma)) \rightarrow I_{Q_1}(\text{St}_P^{Q_1}(\sigma))$. Recall that the kernel of $I_P^Q(\sigma) \ni \varphi \mapsto \varphi(T_{n_{w_Q}w_P}) \in \sigma$ is $\sum_{Q \supset P_1 \supseteq P} I_{P_1}(e_{P_1}(\sigma))$ and hence it gives an identification $\sigma \simeq \text{St}_P^Q(\sigma)$ as vector spaces by Lemma 2.6.

Lemma 3.16. *The homomorphism $I_Q(\text{St}_P^Q(\sigma)) \rightarrow I_{Q_1}(\text{St}_P^{Q_1}(\sigma))$ is given by $\varphi \mapsto (X \mapsto \varphi(XT_{n_{w_{Q_1}w_Q}}))$. (Here we identify $\text{St}_P^Q(\sigma)$ and $\text{St}_P^{Q_1}(\sigma)$ with σ .)*

Proof. Since $I_P^Q(\sigma) \rightarrow \text{St}_P^Q(\sigma)$ is given by $\varphi \mapsto \varphi(T_{n_{w_Q}w_P})$ (under the identification $\sigma = \text{St}_P^Q(\sigma)$), $I_P(\sigma) \rightarrow I_Q(\text{St}_P^Q(\sigma))$ is given by $\varphi \mapsto (X \mapsto \varphi(XT_{n_{w_Q}w_P}))$. Now recall the following commutative diagram which defines the homomorphism in the lemma.

$$\begin{array}{ccc} I_P(\sigma) & \longrightarrow & I_Q(\text{St}_P^Q(\sigma)) \\ \parallel & & \downarrow \\ I_P(\sigma) & \longrightarrow & I_{Q_1}(\text{St}_P^{Q_1}(\sigma)). \end{array}$$

Let $\varphi \in I_Q(\text{St}_P^Q(\sigma))$ and take $\tilde{\varphi} \in I_P(\sigma)$ which is a lift of φ . Then we have $\varphi(X) = \tilde{\varphi}(XT_{n_{w_Q}w_P})$. Let φ' be the image of φ . Then the above commutative diagram we have $\varphi'(X) = \tilde{\varphi}(XT_{n_{w_{Q_1}w_Q}})$. Since $n_{w_{Q_1}w_P} = n_{w_{Q_1}w_Q}n_{w_Qw_P}$, we have $\tilde{\varphi}(XT_{n_{w_{Q_1}w_P}}) = \tilde{\varphi}(XT_{n_{w_{Q_1}w_Q}}T_{n_{w_Qw_P}}) = \varphi(XT_{n_{w_{Q_1}w_Q}})$. $\square$

Proof of Proposition 3.15. Let $\varphi \in I'_Q(e_Q(\sigma))$ and $\varphi' \in I'_{Q_1}(e_{Q_1}(\sigma))$ its image. Then $\varphi \circ \iota \in I_Q(\text{St}_P^Q(\sigma_{\ell-P}^{\iota_P}))$ and $\varphi' \circ \iota \in I_{Q_1}(\text{St}_P^{Q_1}(\sigma_{\ell-P}^{\iota_P}))$. By the above lemma, we have $\varphi' \circ \iota(X) = \varphi \circ \iota(XT_{n_{w_{Q_1}w_Q}})$. Hence $\varphi'(\iota(X)) = \varphi(\iota(XT_{n_{w_{Q_1}w_Q}})) = (-1)^{\ell(w_{Q_1}w_Q)}\varphi(\iota(X)T_{n_{w_{Q_1}w_Q}}^*)$ for any $X \in \mathcal{H}$. Therefore, for any $X \in \mathcal{H}$, we have $\varphi'(X) = (-1)^{\ell(w_{Q_1}w_Q)}\varphi(XT_{n_{w_{Q_1}w_Q}}^*)$ for any $X \in \mathcal{H}$. $\square$

By Proposition 3.15, $\varphi \in \text{Ker}(I'_Q(e_Q(\sigma)) \rightarrow \bigoplus_{Q_1 \supsetneq Q} I'_{Q_1}(e_{Q_1}(\sigma)))$ if and only if $\varphi(T_{n_w}^*) = 0$ for any $w \in \bigcup_{Q_1 \supsetneq Q} W_0^{Q_1}w_{Q_1}w_Q$. We get the following description of the kernel appearing in Lemma 3.9.

Lemma 3.17. *Let $w \in W_0^Q$. We have $\Delta_{wwQ} \neq \Delta \setminus \Delta_Q$ if and only if for some $Q_1 \supsetneq Q$, $w \in W_0^{Q_1} w_{Q_1} w_Q$. Hence we have*

$$\begin{aligned} & \text{Ker} \left(I'_Q(e_Q(\sigma)) \rightarrow \bigoplus_{Q_1 \supsetneq Q} I'_{Q_1}(e_{Q_1}(\sigma)) \right) \\ &= \{ \varphi \in I'_Q(e_Q(\sigma)) \mid \varphi(T_{n_w}^*) = 0 \text{ for any } w \in W_0^Q \text{ such that } \Delta_{wwQ} \neq \Delta \setminus \Delta_Q \}. \end{aligned}$$

Proof. Let $w \in W_0^Q$ and assume that for some $Q_1 \supsetneq Q$ and $v \in W_0^{Q_1}$, we have $w = vw_{Q_1}w_Q$. Then for $\alpha \in \Delta_{Q_1} \setminus \Delta_Q$, $w_{Q_1}(\alpha) \in \Sigma_{Q_1}^-$. Since $v \in W_0^{Q_1}$, we have $vw_{Q_1}(\alpha) < 0$. Hence $ww_Q(\alpha) < 0$. Therefore $\alpha \notin \Delta_{wwQ}$. Hence $\Delta_{wwQ} \neq \Delta \setminus \Delta_Q$.

Assume that $\Delta_{wwQ} \neq \Delta \setminus \Delta_Q$. Since $w \in W_0^Q$, for any $\alpha \in \Delta_Q$ we have $ww_Q(\alpha) < 0$. Hence $\alpha \notin \Delta_{wwQ}$. Therefore $\Delta_Q \subset \Delta \setminus \Delta_{wwQ}$, namely we have $\Delta_{wwQ} \subset \Delta \setminus \Delta_Q$. Hence $\Delta_{wwQ} \not\supset \Delta \setminus \Delta_Q$. Take $\alpha \in (\Delta \setminus \Delta_Q) \setminus \Delta_{wwQ}$. Let Q_1 be a parabolic subgroup corresponding to $\Delta_Q \cup \{\alpha\}$. Then we have $\Delta_{Q_1} \subset \Delta \setminus \Delta_{wwQ}$. If $\beta \in \Delta_{Q_1}$, then $w_{Q_1}(\beta) \in -\Delta_{Q_1} \subset -(\Delta \setminus \Delta_{wwQ})$. Hence $ww_Q w_{Q_1}(\beta) > 0$. Therefore $ww_Q w_{Q_1} \in W_0^{Q_1}$. We have $w \in W_0^{Q_1} w_{Q_1} w_Q$. $\square$

3.6. Complex. In this subsection, we prove that the sequence in Lemma 3.9 is a complex, namely the composition

$$I_Q(e_Q(\sigma)) \rightarrow I'_Q(e_Q(\sigma)) \rightarrow I'_{Q_1}(e_{Q_1}(\sigma))$$

is zero for any parabolic subgroup $Q_1 \supsetneq Q$.

We have the following diagram

$$\begin{array}{ccccc} I_Q(e_Q(\sigma)) & \xrightarrow{\quad\quad\quad} & I'_Q(e_Q(\sigma)) & \xrightarrow{\quad\quad\quad} & I'_{Q_1}(e_{Q_1}(\sigma)) \\ \downarrow \wr & & \downarrow \wr & \nearrow & \\ I_{Q_1} I_Q^{Q_1}(e_Q(\sigma)) & \longrightarrow & I'_{Q_1}(I_Q^{Q_1}(e_Q(\sigma))) & \longrightarrow & I'_{Q_1}(I_Q^{Q_1'}(e_Q(\sigma))). \end{array}$$

This is commutative by Lemma 3.5. The sequence

$$I'_{Q_1}(I_Q^{Q_1}(e_Q(\sigma))) \rightarrow I'_{Q_1}(I_Q^{Q_1'}(e_Q(\sigma))) \rightarrow I'_{Q_1}(e_{Q_1}(\sigma))$$

comes from

$$I_Q^{Q_1}(e_Q(\sigma)) \rightarrow I_Q^{Q_1'}(e_Q(\sigma)) \rightarrow e_{Q_1}(\sigma).$$

Since $R_Q^{Q_1}(e_{Q_1}(\sigma)) = 0$ by [Abe16, Lemma 5.17], this composition is zero.

3.7. Exactness. Now we finish the proof of Lemma 3.9 by proving the following lemma.

Lemma 3.18. *We have*

$$\text{Im} \left(I_Q(e_Q(\sigma)) \rightarrow I'_Q(e_Q(\sigma)) \right) \supset \text{Ker} \left(I'_Q(e_Q(\sigma)) \rightarrow \bigoplus_{Q_1 \supsetneq Q} I'_{Q_1}(e_{Q_1}(\sigma)) \right).$$

We start with the following lemma.

Lemma 3.19. *Let $\psi \in I'_Q(e_Q(\sigma))$ and $w \in W_0^Q$. Then we have $\psi(T_{n_w}^*) = \sum_{v \in W_0^Q, v \leq w} \psi(T_{n_v})$.*

Proof. The same argument as the proof of [AHHV17, IV.9 Proposition] implies

$$\psi(T_{n_w}^*) = \sum_{v \leq w} \psi(T_{n_v}).$$

If $v \notin W_0^Q$, then there exists $v_1 \in W_0^Q$ and $v_2 \in W_{0,Q} \setminus \{1\}$ such that $v = v_1 v_2$. We have $\ell(v_1 v_2) = \ell(v_1) + \ell(v_2)$. Hence $\psi(T_{n_v}) = \psi(T_{n_{v_1}} T_{n_{v_2}}) = \psi(T_{n_{v_1}}) e_Q(\sigma)(T_{n_{v_2}}^Q) = 0$ by the definition of $e_Q(\sigma)$. We get the lemma. $\square$

Let μ^Q be the Möbius function associated to $(W_0^Q, \leq)$. Then by the above lemma, we have

$$\psi(T_{n_w}) = \sum_{v \leq w, v \in W_0^Q} \mu^Q(v, w) \psi(T_{n_v}^*).$$

By [Deo77, Theorem 1.2], μ^Q is given by

$$\mu^Q(v, w) = \begin{cases} 0 & (\text{there exists } \alpha \in \Delta_Q \text{ such that } v s_\alpha \leq w), \\ (-1)^{\ell(v) + \ell(w)} & (\text{otherwise}). \end{cases}$$

Now we assume that $\psi \in \text{Ker}(I'_Q(e_Q(\sigma)) \rightarrow \bigoplus_{Q_1 \supsetneq Q} I'_{Q_1}(e_{Q_1}(\sigma)))$. Then we have $\psi(T_{n_w}^*) = 0$ if $\Delta_{ww_Q} \neq \Delta \setminus \Delta_Q$ by Lemma 3.17. Hence we have $\psi(T_{n_w}) = \sum_{v \leq w, v \in W_0^Q, \Delta_{vw_Q} = \Delta \setminus \Delta_Q} \mu^Q(v, w) \psi(T_{n_v}^*)$. Let w_c be the longest element of the finite Weyl group of the parabolic subgroup corresponding to $\Delta \setminus \Delta_Q$. Then $\Delta_{w_G w_c} = \Delta \setminus \Delta_Q$ and $w = w_G w_c$ is maximal in $\{w \in W_0 \mid \Delta_w = \Delta \setminus \Delta_Q\}$ [Abe, Remark 3.16].

Lemma 3.20. *Let $w \in W_0^Q$ such that $\Delta_{ww_Q} = \Delta \setminus \Delta_Q$. Then we have*

$$\mu^Q(w, w_G w_Q) = \begin{cases} (-1)^{\ell(w_c)} & (w = w_G w_c w_Q), \\ 0 & (w \neq w_G w_c w_Q). \end{cases}$$

We prove this lemma by backward induction on the length of w . For the inductive step, we use the following.

Lemma 3.21. *Let $w \in W_0^Q$ such that $\Delta_{ww_Q} = \Delta \setminus \Delta_Q$, $w \neq w_G w_c w_Q$ and $\alpha \in \Delta$ such that $s_\alpha w w_Q > w w_Q$ and $\Delta_{ww_Q} = \Delta_{s_\alpha w w_Q}$. (Such α exists [Abe, Lemma 3.15].) We have*

- (1) $s_\alpha w \in W_0^Q$.
- (2) $s_\alpha w > w$.

Proof. If $\beta \in \Delta_Q$, then $w_Q(\beta) \in -\Delta_Q$. Hence $s_\alpha w w_Q(w_Q(\beta)) > 0$ since $\Delta_{s_\alpha w w_Q} = \Delta \setminus \Delta_Q$. Therefore $s_\alpha w(\beta) > 0$ for any $\beta \in \Delta_Q$. Namely we have $s_\alpha w \in W_0^Q$. Therefore we have $s_\alpha w, w \in W_0^Q$, $w_Q \in W_{0,Q}$ and $s_\alpha w w_Q > w w_Q$. By [Deo77, Lemma 3.5], $s_\alpha w > w$. $\square$

Proof of Lemma 3.20. Assume that $w = w_G w_c w_Q$ and there exists $\alpha \in \Delta_Q$ such that $ws \leq w_G w_Q$ where $s = s_\alpha$. Then we have $w_G w_c w_Q s \leq w_G w_Q$. Hence $w_c w_Q s \geq w_Q$. Let Q_0 be the parabolic subgroup corresponding to $\Delta \setminus \Delta_Q$. Then $w_c \in W_{0,Q_0}$ and $W_{0,Q} \subset {}^{Q_0}W_0$. Therefore $w_Q, w_Q s \in {}^{Q_0}W_0$. Hence by [Deo77, Lemma 3.5], we have $w_Q s \geq w_Q$. This is a contradiction. Hence $\mu^Q(w, w_G w_Q) = (-1)^{\ell(w) + \ell(w_G w_Q)} = (-1)^{\ell(w_c)}$.

If $w \neq w_G w_c w_Q$, then $w w_Q \neq w_G w_c$. Take α as in the previous lemma. Assume that $s_\alpha w = w_G w_c w_Q$. Since $s_\alpha w_G w_c = w w_Q < s_\alpha w w_Q = w_G w_c$, we

have $(w_G w_c)^{-1}(\alpha) < 0$. Put $\alpha' = -w_G^{-1}(\alpha) \in \Delta$. Then we have $w_c^{-1}(\alpha') > 0$. Hence $\alpha' \in \Delta_Q$. Put $\beta = -w_Q(\alpha') \in \Delta_Q$ and we prove $ws_\beta \leq w_G w_Q$. We have $ws_\beta = s_\alpha w_G w_c w_Q s_\beta = w_G s_{\alpha'} w_c w_Q s_\beta$. Hence it is sufficient to prove that $s_{\alpha'} w_c w_Q s_\beta \geq w_Q$.

We have $\Delta_{w_G s_{\alpha'} w_c} = \Delta_{s_\alpha w_G w_c} = \Delta_{w w_Q} = \Delta \setminus \Delta_Q$. Hence $\Delta_{s_{\alpha'} w_c} = \Delta_Q$. In particular, $s_{\alpha'} w_c \in W_0^Q$. Hence $\ell(s_{\alpha'} w_c w_Q s_\beta) = \ell(s_{\alpha'} w_c) + \ell(w_Q s_\beta)$ as $w_Q, s_\beta \in W_{0,Q}$. Since $\alpha' \in \Delta_Q$, $s_{\alpha'} \in W_{0,Q} \subset W_0^{Q_0}$. Hence we have $\ell(s_{\alpha'} w_c) = \ell(s_{\alpha'}) + \ell(w_c)$. Therefore we get

$$\ell(s_{\alpha'} w_c w_Q s_\beta) = \ell(s_{\alpha'}) + \ell(w_c) + \ell(w_Q s_\beta).$$

Hence $s_{\alpha'} w_c w_Q s_\beta \geq s_{\alpha'} w_Q s_\beta = w_Q$.

Finally assume that $s_\alpha w \neq w_G w_c$ and we prove the lemma by backward induction on $\ell(w w_Q)$. By inductive hypothesis, there exists $\beta \in \Delta_Q$ such that $s_\alpha w s_\beta \leq w_G w_Q$. Since $s_\alpha w \in W_0^Q$, we have $s_\alpha w s_\beta > s_\alpha w$. Therefore we have $s_\alpha w s_\beta > s_\alpha w > w$. By Property $Z(ws_\beta, s_\alpha w s_\beta, s_\beta)$ [Deo77, §1, Remarks (2)], we have $ws_\beta \leq s_\alpha w s_\beta$. Since we have $s_\alpha w s_\beta \leq w_G w_Q$, we get $ws_\beta \leq w_G w_Q$. $\square$

Therefore we get

Lemma 3.22. *If $\psi \in \text{Ker}(I'_Q(e_Q(\sigma)) \rightarrow \bigoplus_{Q_1 \supseteq Q} I'_{Q_1}(e_{Q_1}(\sigma)))$, then $\psi(T_{n_{w_G w_Q}}) = (-1)^{\ell(w_c)} \psi(T_{n_{w_G w_c w_Q}}^*)$.*

Since the image of $I_Q(e_Q(\sigma)) \rightarrow I'_Q(e_Q(\sigma))$ is contained in the kernel by the previous subsection, we get the following lemma.

Lemma 3.23. *Consider a linear map $I'_Q(e_Q(\sigma)) \rightarrow \sigma$ defined by $\psi \mapsto \psi(T_{n_{w_G w_Q}}^*)$. Then the composition $I_Q(e_Q(\sigma)) \rightarrow I'_Q(e_Q(\sigma)) \rightarrow \sigma$ is surjective.*

Proof. Let $\psi \in I'_Q(e_Q(\sigma))$ be the image of $\varphi \in I_Q(e_Q(\sigma))$. Then the characterization of the homomorphism (Proposition 3.1) gives $\varphi(T_{n_{w_G w_Q}}) = \psi(T_{n_{w_G w_Q}})$. The lemma follows from the previous lemma and the surjectivity of the map $I_Q(e_Q(\sigma)) \ni \varphi \mapsto \varphi(T_{n_{w_G w_Q}}) \in \sigma$. $\square$

The following lemma ends the proof of Lemma 3.18, hence that of Theorem 3.6.

Lemma 3.24. *For $w \in W_0^Q$ such that $\Delta_{w w_Q} = \Delta \setminus \Delta_Q$ and $x \in \sigma$, there exists $\psi \in \text{Im}(I_Q(e_Q(\sigma)) \rightarrow I'_Q(e_Q(\sigma)))$ such that for any $v \in W_0^Q$ we have*

$$\psi(T_{n_v}^*) = \begin{cases} x & (v = w), \\ 0 & (v \neq w). \end{cases}$$

We need one lemma.

Lemma 3.25. *Let $w \in W_0^P$ and $\lambda \in \Lambda(1)$. Then for $\varphi \in I'_P(\sigma)$, we have*

$$(\varphi E_{o_+}(\lambda))(T_{n_w}^*) = \begin{cases} \varphi(T_{n_w}^*) \sigma(E_{o_+, P}^P(n_w^{-1} \cdot \lambda)) & (n_w^{-1} \cdot \lambda \in W_P^-(\lambda)), \\ 0 & (n_w^{-1} \cdot \lambda \notin W_P^-(\lambda)). \end{cases}$$

Proof. Set $\varphi^\iota = \varphi \circ \iota$ and $\sigma' = \sigma_{\ell - \ell_P}^{\iota_P}$. Then we have $\varphi^\iota \in I_P(\sigma')$ [Abe16, Proposition 4.11]. Hence, by Proposition 2.4, we have

$$\varphi^\iota(E_{o_-}(\lambda) T_{n_w}) = \begin{cases} \varphi^\iota(T_{n_w}) \sigma'(E_{o_-}^P(n_w^{-1} \cdot \lambda)) & (n_w^{-1} \cdot \lambda \in W_P^-(\lambda)), \\ 0 & (n_w^{-1} \cdot \lambda \notin W_P^-(\lambda)). \end{cases}$$

The left hand side is

$$\varphi'(E_{o_-}(\lambda)T_{n_w}) = \varphi(\iota(E_{o_-}(\lambda))\iota(T_{n_w})) = \varphi((-1)^{\ell(\lambda)}E_{o_+}(\lambda)(-1)^{\ell(n_w)}T_{n_w}^*)$$

by [Vig16, Lemma 5.31]. Therefore if $n_w^{-1} \cdot \lambda \notin W_P^-(\lambda)$, then $\varphi(E_{o_+}(\lambda)T_{n_w}^*) = 0$.

If $n_w^{-1} \cdot \lambda \in W_P^-(\lambda)$, then

$$\begin{aligned} \sigma'(E_{o_-}^P(n_w^{-1} \cdot \lambda)) &= (-1)^{\ell(n_w^{-1} \cdot \lambda) - \ell_P(n_w^{-1} \cdot \lambda)} \sigma(\iota_P(E_{o_-}^P(n_w^{-1} \cdot \lambda))) \\ &= (-1)^{\ell(n_w^{-1} \cdot \lambda)} \sigma(E_{o_+}^P(n_w^{-1} \cdot \lambda)) \end{aligned}$$

again by [Vig16, Lemma 5.31] and [Abe16, Lemma 4.5]. We also have $\varphi'(T_{n_w}) = (-1)^{\ell(n_w)}\varphi(T_{n_w}^*)$. Hence we get

$$\varphi((-1)^{\ell(\lambda)}E_{o_+}(\lambda)(-1)^{\ell(n_w)}T_{n_w}^*) = (-1)^{\ell(n_w)}(-1)^{\ell(n_w^{-1} \cdot \lambda)}\varphi(T_{n_w}^*)\sigma(E_{o_+}^P(n_w^{-1} \cdot \lambda))$$

Since $\ell(n_w^{-1} \cdot \lambda) = \ell(\lambda)$ [Abe16, Lemma 2.15], we get the lemma. $\square$

Proof of Lemma 3.24. We prove the lemma by induction on the length of ww_Q . Assume that $ww_Q = w_Gw_c$, namely $w = w_Gw_cw_Q$. Let $\lambda_P^- \in Z(W_P(1))$ as in Proposition 2.2. Notice that $n_{w_Q}^{-1} \cdot \lambda_P^-$ is P -negative since w_Q (in fact, any element in W_0) preserves $\Sigma^+ \setminus \Sigma_P^+ = \Sigma_{P_2}^+$. We also have that $n_{w_Q}^{-1} \cdot \lambda_P^- \in Z(W_P(1))$ since n_{w_Q} normalizes $W_P(1)$. Hence $e_Q(\sigma)(T_{n_{w_Q}^{-1} \cdot \lambda_P^-}^{Q*}) = \sigma(T_{n_{w_Q}^{-1} \cdot \lambda_P^-}^{P*})$ is invertible. Take $\psi_0 \in \text{Im}(I_Q(e_Q(\sigma)) \rightarrow I'_Q(e_Q(\sigma)))$ such that $\psi_0(T_{n_{w_Gw_cw_Q}}^*) = xe_Q(\sigma)(T_{n_{w_Q}^{-1} \cdot \lambda_P^-}^{Q*})^{-1}$.

Put $\lambda = n_{w_Gw_c} \cdot \lambda_P^-$ and set $\psi = \psi_0 E_{o_+}(\lambda)$. Let $v \in W_0^Q$. If $v \neq w_Gw_cw_Q$, then since v and $w_Gw_cw_Q$ are in W_0^Q , we have $v \notin w_Gw_cw_Q W_{0,Q}$. Hence $(w_Gw_c)^{-1}v \notin W_{0,Q}$. Therefore there exists $\alpha \in \Sigma^+ \setminus \Sigma_Q^+$ such that $(w_Gw_c)^{-1}v(\alpha) < 0$. Since $\Sigma^+ \setminus \Sigma_Q^+ \subset \Sigma_{P_2}$ and Σ_{P_2} is stabilized by W_0 , we have $(w_Gw_c)^{-1}v(\alpha) \in \Sigma_{P_2}^- = \Sigma^- \setminus \Sigma_P^-$. Hence $\langle (w_Gw_c)^{-1}v(\alpha), \nu(\lambda_P^-) \rangle < 0$. The left hand side is $\langle \alpha, \nu(n_v^{-1} \cdot \lambda) \rangle$. Hence $n_v^{-1} \cdot \lambda$ is not Q -negative. Therefore $\psi(T_{n_v}) = (\psi_0 E_{o_+}(\lambda))(T_{n_v}) = 0$ if $v \neq w_Gw_cw_Q$ by the above lemma.

Assume that $v = w_Gw_cw_Q$. Then $n_v^{-1} \cdot \lambda = n_{w_Q}^{-1} \cdot \lambda_P^-$. Since λ_P^- is dominant, it is Q -negative. The set of Q -negative elements is stable under the conjugate action of $W_Q(1)$. Hence $n_{w_Q}^{-1} \cdot \lambda_P^-$ is also Q -negative. Therefore we have $\psi(T_{n_{w_Gw_cw_Q}}^*) = (\psi_0 E_{o_+}(\lambda))(T_{n_{w_Gw_cw_Q}}^*) = \psi_0(T_{n_{w_Gw_cw_Q}}^*)e_Q(\sigma)(T_{n_{w_Q}^{-1} \cdot \lambda_P^-}^{Q*}) = x$ by the previous lemma. We get the lemma when $w = w_Gw_cw_Q$.

Assume that $w \neq w_Gw_cw_Q$ and take α such that $s_\alpha ww_Q > ww_Q$ and $\Delta_{s_\alpha ww_Q} = \Delta_{ww_Q}$ as in Lemma 3.21. Then $s_\alpha w > w$ by Lemma 3.21. Set $s = s_\alpha$. By inductive hypothesis, there exists $\psi_0 \in \text{Im}(I_Q(e_Q(\sigma)) \rightarrow I'_Q(e_Q(\sigma)))$ such that for $v \in W_0^Q \setminus \{s_\alpha w\}$, $\psi_0(T_{n_v}^*) = 0$ and $\psi_0(T_{n_{s_\alpha w}}^*) = x$. We prove that $\psi = \psi_0 T_{n_s}$ satisfies the condition of the lemma. First we have

$$\psi(T_{n_w}^*) = \psi_0(T_{n_s} T_{n_w}^*) = \psi_0((T_{n_s}^* - c_{n_s})T_{n_w}^*) = \psi_0(T_{n_s}^* T_{n_w}^*) - \psi_0(c_{n_s} T_{n_w}^*).$$

Since $sw > w$, we have $T_{n_s}^* T_{n_w}^* = T_{n_{sw}}^*$. Hence $\psi_0(T_{n_s}^* T_{n_w}^*) = x$. Since $\psi_0(T_{n_w}^*) = 0$, we have $\psi_0(c_{n_s} T_{n_w}^*) = \psi_0(T_{n_s}^*)e_Q(\sigma)(n_w^{-1} \cdot c_{n_s}) = 0$. Hence $\psi(T_{n_w}^*) = x$.

Assume that $v \neq w$. If $\ell(s) + \ell(v) > \ell(sv)$, then $T_{n_s} T_{n_v}^* = T_{n_s} T_{n_s}^* T_{n_v}^* = 0$. Hence $\psi(T_{n_v}^*) = 0$. If $\ell(s) + \ell(v) = \ell(sv)$, then

$$\psi(T_{n_v}^*) = \psi_0(T_{n_s} T_{n_v}^*) = \psi_0((T_{n_s}^* - c_{n_s})T_{n_v}^*) = \psi_0(T_{n_s}^* T_{n_v}^*) - \psi_0(c_{n_s} T_{n_v}^*).$$

Since $v \neq w$, $sv \neq sw$. Hence $\psi_0(T_{n_{sv}}^*) = 0$. Since $sv > v$ and $s(sw) < sw$, we have $v \neq sw$. Hence $\psi_0(c_{n_s} T_{n_v}^*) = \psi_0(T_{n_v}^*) e_Q(\sigma)(n_v^{-1} \cdot c_{n_s}) = 0$. Therefore we get $\psi(T_{n_v}^*) = 0$. $\square$

3.8. Supersingular modules.

Proposition 3.26. *If π is supersingular, then π^ι is also supersingular.*

Proof. For each $W(1)$ -orbit $\mathcal{O} \subset \Lambda(1)$ such that $\ell(\mathcal{O}) > 0$, we have

$$\iota(z_{\mathcal{O}}) = \sum_{\lambda \in \mathcal{O}} \iota(E_{o_-}(\lambda)) = \sum_{\lambda \in \mathcal{O}} (-1)^{\ell(\mathcal{O})} E_{o_+}(\lambda) = (-1)^{\ell(\mathcal{O})} z_{\mathcal{O}}$$

by [Vig16, Lemma 5.31]. Hence $\pi(z_{\mathcal{O}}^n) = 0$ implies $\pi^\iota(z_{\mathcal{O}}^n) = 0$. $\square$

Proposition 3.27. *Assume that C is a field. Let $\pi = \pi_{\chi, J, V}$ be a simple supersingular representation. Then $\pi^\iota = \pi_{\chi, S_{\text{aff}, \chi} \setminus J, V}$.*

Proof. Let $\Xi = \Xi_{\chi, J}$ be the character of $\mathcal{H}_{\text{aff}}$ parametrized by (χ, J) . The representation π is given by $\pi = (V \otimes \Xi_{\chi, J}) \otimes_{\mathcal{H}_{\text{aff}} C[\Omega(1)_{\Xi}]} \mathcal{H}$. The homomorphism ι preserves $\mathcal{H}_{\text{aff}}$ and $C[\Omega(1)_{\Xi}]$. (On $C[\Omega(1)]$, ι is identity.) Hence we get $\pi^\iota = (V^\iota \otimes \Xi^\iota) \otimes_{\mathcal{H}_{\text{aff}} C[\Omega(1)_{\Xi}]} \mathcal{H}$. Since ι is trivial on $C[\Omega(1)_{\Xi}]$, $V^\iota = V$. Let (χ', J') be the pair such that Ξ^ι is parametrized by (χ', J') . The character χ' is a direct summand of $V^\iota|_{Z_\kappa \cap W_{\text{aff}}(1)}$ and since $V^\iota = V$, we have $V^\iota|_{Z_\kappa \cap W_{\text{aff}}(1)} = V|_{Z_\kappa \cap W_{\text{aff}}(1)}$. Since $V|_{Z_\kappa \cap W_{\text{aff}}(1)}$ is a direct sum of χ , $\chi' = \chi$. The subset $J \subset S_{\text{aff}, \chi}$ satisfies

$$\Xi(T_s^\sim) = \begin{cases} 0 & (s \in J), \\ \chi(c_s^\sim) & (s \in S_{\text{aff}, \chi} \setminus J). \end{cases}$$

where $\tilde{s} \in W_{\text{aff}}(1)$ is a lift of s . We have $\Xi(\iota(T_s^\sim)) = -\Xi(T_s^\sim - c_s^\sim) = -\Xi(T_s^\sim) + \chi(c_s^\sim)$. Therefore we have

$$\Xi^\iota(T_s^\sim) = \begin{cases} \chi(c_s^\sim) & (s \in J), \\ 0 & (s \in S_{\text{aff}, \chi} \setminus J). \end{cases}$$

We have $J' = \{s \in S_{\text{aff}, \chi} \mid \Xi^\iota(T_{n_s}) = 0\}$. Hence $J' = S_{\text{aff}, \chi} \setminus J$. $\square$

3.9. Simple modules. Assume that C is an algebraically closed field. Summarizing the results in this section, we have the following. We need notation. By [Abe16, Remark 4.6], $T_w^P \mapsto (-1)^{\ell(w) - \ell_P(w)} T_w^P$ is an algebra homomorphism of $\mathcal{H}_P$. This preserves the subalgebra $C[\Omega_P(1)]$. Let Ξ be a character of $\mathcal{H}_{\text{aff}, P}$. Then the above homomorphism also preserves $C[\Omega_P(1)_{\Xi}]$ since the homomorphism is trivial on $\mathcal{H}_{\text{aff}, P}$ by [Abe16, Lemma 4.7]. For a $C[\Omega_P(1)_{\Xi}]$ -module V , let $V_{\ell - \ell_P}$ be the pull-back of V by this homomorphism.

Theorem 3.28. *Let $I(P; \chi, J, V; Q)$ be a simple representation. Then we have $I(P; \chi, J, V; Q)^\iota = I(P; \chi, S_{\text{aff}, P, \chi} \setminus J, V_{\ell - \ell_P}; Q^c)$ where $\Delta_{Q^c} = \Delta_P \cup (\Delta(\sigma) \setminus \Delta_Q)$.*

Proof. Since $I(P; \chi, J, V; Q) = I_{P(\sigma)}(\text{St}_Q^{P(\sigma)}(\pi_{\chi, J, V}))$ is simple, by Corollary 3.3, we have $I(P; \chi, J, V; Q)^\iota = I_{P(\sigma)}((\text{St}_Q^{P(\sigma)}(\pi_{\chi, J, V}))_{\ell - \ell_P}^{\iota_{P(\sigma)}})$. By Theorem 3.6, Proposition 3.27 and [Abe16, Lemma 4.9], we have

$$\begin{aligned} I(P; \chi, J, V; Q)^\iota &\simeq I_{P(\sigma)}(\text{St}_{Q^c}^{P(\sigma)}((\pi_{\chi, J, V})_{\ell - \ell_P}^{\iota_P})) \\ &\simeq I_{P(\sigma)}(\text{St}_{Q^c}^{P(\sigma)}((\pi_{\chi, S_{\text{aff}, P, \chi} \setminus J, V})_{\ell - \ell_P})). \end{aligned}$$

Let Ξ be a character of $\mathcal{H}_{\text{aff},P}$ defined by the pair χ and $S_{\text{aff},P,\chi} \setminus J$. Put $\mathcal{H}_{P,\Xi} = \mathcal{H}_{\text{aff},P} C[\Omega_P(1)\Xi]$. Then $\pi_{\chi, S_{\text{aff},P,\chi} \setminus J, V} = (\Xi \otimes V) \otimes_{\mathcal{H}_{P,\Xi}} \mathcal{H}_P$. Let $f: \mathcal{H}_P \rightarrow \mathcal{H}_P$ be an algebra homomorphism defined by $f(T_w^P) = (-1)^{\ell(w) - \ell_P(w)} T_w^P$. Then f preserves $\mathcal{H}_{\text{aff},P}$ and $C[\Omega_P(1)\Xi]$ and we have $(\pi_{\chi, S_{\text{aff},P,\chi} \setminus J, V})_{\ell - \ell_P} = \pi_{\chi, S_{\text{aff},P,\chi} \setminus J, V} \circ f = ((\Xi \circ f) \otimes (V \circ f)) \otimes_{\mathcal{H}_{P,\Xi}} \mathcal{H}_P$. By the definition, $V \circ f = V_{\ell - \ell_P}$. By [Abe16, Lemma 4.7], f is identity on $\mathcal{H}_{\text{aff},P}$. Hence $\Xi \circ f = \Xi$. Hence $(\pi_{\chi, S_{\text{aff},P,\chi} \setminus J, V})_{\ell - \ell_P} = (\Xi \otimes V_{\ell - \ell_P}) \otimes_{\mathcal{H}_{P,\Xi}} \mathcal{H}$ and we get the theorem. $\square$

3.10. Structure of I'_P . Assume that C is an algebraically closed field.

Proposition 3.29. *Let P be a parabolic subgroup and σ a simple supersingular representation of $\mathcal{H}_P$. Then for each parabolic subgroup Q between P and $P(\sigma)$, there exists a submodule $\pi_Q \subset I'_P(\sigma)$ such that*

- (1) if $Q_1 \subset Q_2$ then $\pi_{Q_1} \subset \pi_{Q_2}$.
- (2) $\pi_Q / \sum_{Q_1 \subsetneq Q} \pi_{Q_1} = I(P, \sigma, Q)$.

Compare with $I_Q(e_Q(\sigma)) \subset I_P(\sigma)$. In other words, the structure of $I'_P(\sigma)$ is “opposite to” that of $I_P(\sigma)$.

Proof. First assume that $P(\sigma) = G$. Put $\sigma' = \sigma_{\ell - \ell_P}^{\ell_P}$. Then $I'_P(\sigma) = I_P(\sigma')^\iota$. Set $\pi_Q = I_{Q^c}(e_{Q^c}(\sigma'))^\iota \subset I'_P(\sigma)$ where $\Delta_{Q^c} = (\Delta \setminus \Delta_Q) \cup \Delta_P$. Then the first condition is satisfied. Since $Q_1 \subset Q_2$ if and only if $Q_1^c \supset Q_2^c$, we have

$$\begin{aligned} \pi_Q / \sum_{Q_1 \subsetneq Q} \pi_{Q_1} &= \left(I_{Q^c}(e_{Q^c}(\sigma')) / \sum_{Q_1 \subsetneq Q} I_{Q_1^c}(e_{Q_1^c}(\sigma')) \right)^\iota \\ &= \left(I_{Q^c}(e_{Q^c}(\sigma')) / \sum_{Q_1 \supsetneq Q^c} I_{Q_1}(e_{Q_1}(\sigma')) \right)^\iota \\ &= (\text{St}_{Q^c}(\sigma'))^\iota = \text{St}_Q(\sigma) \end{aligned}$$

by Theorem 3.6. By the assumption $P(\sigma) = G$, we have $\text{St}_Q(\sigma) = I(P, \sigma, Q)$. We get the proposition in this case.

In general, applying the proposition for $I_P^{P(\sigma)'}(\sigma)$, we get $\pi'_Q \subset I_P^{P(\sigma)'}(\sigma)$ for each $P(\sigma) \supset Q \supset P$. Put $\pi_Q = I'_{P(\sigma)}(\pi'_Q)$. The first condition is obvious. For the second condition, we have

$$\begin{aligned} \pi_Q / \sum_{Q_1 \subsetneq Q} \pi_{Q_1} &= I'_{P(\sigma)}(\pi'_Q) / \sum_{Q_1 \subsetneq Q} I'_{P(\sigma)}(\pi'_{Q_1}) \\ &\simeq I'_{P(\sigma)} \left(\pi'_Q / \sum_{Q_1 \subsetneq Q} \pi'_{Q_1} \right) \\ &\simeq I'_{P(\sigma)}(\text{St}_Q^{P(\sigma)}(\sigma)). \end{aligned}$$

Since $I(P, \sigma, Q) = I_{P(\sigma)}(\text{St}_Q^{P(\sigma)}(\sigma))$ is simple, by Corollary 3.3, we have $I(P, \sigma, Q) \simeq I'_{P(\sigma)}(\text{St}_Q^{P(\sigma)}(\sigma))$. Now we get the proposition. $\square$

4. DUAL

We have an anti-automorphism $\zeta = \zeta_G: \mathcal{H} \rightarrow \mathcal{H}$ defined by $\zeta(T_w) = T_{w^{-1}}$. Hence for a representation π , its linear dual $\pi^* = \text{Hom}_C(\pi, C)$ has a structure of

a right $\mathcal{H}$ -module defined by $(fX)(v) = f(v\zeta(X))$ for $f \in \pi^*$, $v \in \pi$ and $X \in \mathcal{H}$. Since any simple representation is finite-dimensional, if π is simple then π^* is again simple. In this section, we compute π^* .

Lemma 4.1. *We have $\zeta(T_w^*) = T_{w^{-1}}^*$.*

Proof. In $\mathcal{H}[q_s^{\pm 1}]$, we have $\zeta(T_w^*) = \zeta(q_w T_{w^{-1}}^{-1}) = q_w T_w^{-1} = T_{w^{-1}}^*$. $\square$

4.1. Parabolic inductions. In this subsection, we calculate $I_P(\sigma)^*$. Let $P' = n_{w_G w_P} P^{\text{op}} n_{w_G w_P}^{-1}$. Then we have $I_P(\sigma) \simeq n_{w_G w_P} \sigma \otimes_{(\mathcal{H}_{P'}^+, j_{P'}^+)} \mathcal{H}$ by [Abe16, Proposition 2.21]. Hence we have

$$\begin{aligned} \text{Hom}_C(I_P(\sigma), C) &\simeq \text{Hom}_C(n_{w_G w_P} \sigma \otimes_{(\mathcal{H}_{P'}^+, j_{P'}^+)} \mathcal{H}, C) \\ &\simeq \text{Hom}_{(\mathcal{H}_{P'}^+, j_{P'}^+)}(\mathcal{H}, \text{Hom}_C(n_{w_G w_P} \sigma, C)). \end{aligned}$$

Therefore $I_P(\sigma)^* \simeq \text{Hom}_{(\mathcal{H}_{P'}^+, j_{P'}^+)}(\mathcal{H}, \text{Hom}_C(n_{w_G w_P} \sigma, C))$ here the action on the right hand side is twisted by ζ . Let $\varphi \in \text{Hom}_{(\mathcal{H}_{P'}^+, j_{P'}^+)}(\mathcal{H}, \text{Hom}_C(n_{w_G w_P} \sigma, C))$ and set $\varphi^\zeta = \varphi \circ \zeta$. Let $w \in W_{P'}(1)$ which is P' -negative. Then w^{-1} is P' -positive. Hence for $X \in \mathcal{H}$ and $x \in n_{w_G w_P} \sigma$, we have $\varphi^\zeta(X T_w)(x) = \varphi(\zeta(X T_w))(x) = \varphi(T_{w^{-1}} \zeta(X))(x) = (T_{w^{-1}}^{P'} \varphi(\zeta(X)))(x) = \varphi(\zeta(X))(x T_w^{P'}) = \varphi^\zeta(X)(x \zeta_P(T_w^{P'}))$. (Here we regard $\text{Hom}_C(n_{w_G w_P} \sigma, C)$ as a left $\mathcal{H}_{P'}$ -module.) Therefore

$$\varphi^\zeta \in \text{Hom}_{(\mathcal{H}_{P'}^-, j_{P'}^-)}(\mathcal{H}, (n_{w_G w_P} \sigma)^*).$$

For $X, Y \in \mathcal{H}$, we have $(\varphi Y)^\zeta(X) = (\varphi Y)(\zeta(X)) = \varphi(\zeta(X) \zeta(Y)) = \varphi(\zeta(YX)) = \varphi^\zeta(YX) = (\varphi^\zeta Y)(X)$. Hence $\varphi \mapsto \varphi^\zeta$ induces

$$I_P(\sigma)^* \simeq \text{Hom}_{(\mathcal{H}_{P'}^-, j_{P'}^-)}(\mathcal{H}, (n_{w_G w_P} \sigma)^*) = I_{P'}'(n_{w_G w_P} \sigma^*).$$

Proposition 4.2. *We have $I_P(\sigma)^* \simeq I_{P'}'(n_{w_G w_P} \sigma^*)$.*

The same calculation shows that

Proposition 4.3. *We have $I_P'(\sigma)^* \simeq I_P'(n_{w_G w_P} \sigma^*)$.*

Remark 4.4. These propositions are true for any commutative ring C .

4.2. Steinberg modules. Let P be a parabolic subgroup, σ an $\mathcal{H}_P$ -module such that $P(\sigma) = G$ and P_2 the parabolic subgroup corresponding to $\Delta \setminus \Delta_P$. We calculate $(\text{St}_Q \sigma)^*$.

Proposition 4.5. *Let Q be a parabolic subgroup containing P and put $Q' = n_{w_G w_Q} Q^{\text{op}} n_{w_G w_Q}^{-1}$. Then $(\text{St}_Q \sigma)^* \simeq \text{St}_{Q'} \sigma^*$.*

We start with the case of $Q = G$.

Lemma 4.6. *We have $e_G(\sigma)^* \simeq e_G(\sigma^*)$.*

Proof. Let $f \in e_G(\sigma)^*$ and $x \in e_G(\sigma)$. For $w \in W_P(1)$, we have

$$\begin{aligned} (f e_G(\sigma)^*(T_w^*))(x) &= f(x e_G(\sigma)(\zeta(T_w^*))) = f(x e_G(\sigma)(T_{w^{-1}}^*)) \\ &= f(x \sigma(T_{w^{-1}}^{P*})) = f(x \sigma(\zeta_P(T_w^{P*}))) = (f \sigma^*(T_w^{P*}))(x). \end{aligned}$$

Hence $e_G(\sigma)^*(T_w^*) = \sigma^*(T_w^{P*})$. For $w \in W_{\text{aff}, P_2}(1)$, we have $(f e_G(\sigma)^*(T_w^*))(x) = f(x e_G(\sigma)(\zeta(T_w^*))) = f(x e_G(\sigma)(T_{w^{-1}}^*)) = f(x)$. Hence $e_G(\sigma)^*(T_w^*) = 1$. Therefore by the characterization of $e_G(\sigma^*)$, we have the lemma. $\square$

Proof of Proposition 4.5. By Lemma 3.9, we have the following exact sequence

$$0 \rightarrow \mathrm{St}_Q(\sigma) \rightarrow I'_Q(e_Q(\sigma)) \rightarrow \bigoplus_{Q_1 \supsetneq Q} I'_{Q_1}(e_{Q_1}(\sigma)).$$

Taking dual, we get an exact sequence

$$\bigoplus_{Q_1 \supsetneq Q} I'_{Q_1}(e_{Q_1}(\sigma))^* \rightarrow I'_Q(e_Q(\sigma))^* \rightarrow \mathrm{St}_Q(\sigma)^* \rightarrow 0.$$

Put $Q' = n_{w_G w_Q} Q^{\mathrm{op}} n_{w_G w_Q}^{-1}$. Then by Proposition 4.3, we have $I'_Q(e_Q(\sigma))^* = I_{Q'}(n_{w_G w_Q} e_Q(\sigma)^*) = I_{Q'}(e_{Q'}(\sigma^*)) = I_{Q'}(e_{Q'}(\sigma^*))$ by Lemma 4.6 and [Abe16, Lemma 2.27]. Put $Q'_1 = n_{w_G w_{Q_1}} Q_1^{\mathrm{op}} n_{w_G w_{Q_1}}^{-1}$. Then

$$\bigoplus_{Q_1 \supsetneq Q} I_{Q'_1}(e_{Q'_1}(\sigma^*)) \rightarrow I_{Q'}(e_{Q'}(\sigma^*)) \rightarrow \mathrm{St}_Q(\sigma)^* \rightarrow 0.$$

Since $\Delta_{Q'} = -w_G(\Delta_Q)$ and $Q'_1 = -w_G(\Delta_{Q_1})$, we have $Q_1 \supsetneq Q$ if and only if $Q'_1 \supsetneq Q'$. Hence

$$\bigoplus_{Q_2 \supsetneq Q'} I_{Q_2}(e_{Q_2}(\sigma^*)) \rightarrow I_{Q'}(e_{Q'}(\sigma^*)) \rightarrow \mathrm{St}_Q(\sigma)^* \rightarrow 0.$$

By the lemma below, we get $\mathrm{St}_Q(\sigma)^* = \mathrm{St}_Q(\sigma^*)$. $\square$

Lemma 4.7. *Let $Q_1 \supset Q$ be parabolic subgroups. Put $Q' = n_{w_G w_Q} Q^{\mathrm{op}} n_{w_G w_Q}^{-1}$ and $Q'_1 = n_{w_G w_{Q_1}} Q_1^{\mathrm{op}} n_{w_G w_{Q_1}}^{-1}$. Then the homomorphism induced by $I'_Q(e_Q(\sigma)) \rightarrow I'_{Q_1}(e_{Q_1}(\sigma))$ with the dual is the inclusion $I_{Q'_1}(e_{Q'_1}(\sigma^*)) \hookrightarrow I_{Q'}(e_{Q'}(\sigma^*))$ times $(-1)^{\ell(w_{Q_1} w_Q)}$.*

Proof. By Proposition 3.15, the homomorphism $I'_Q(e_Q(\sigma)) \rightarrow I'_{Q_1}(e_{Q_1}(\sigma))$ is given by $\varphi \mapsto (X \mapsto (-1)^{\ell(w_{Q_1} w_Q)} \varphi(X T_{n_{w_{Q_1} w_Q}}^*))$. We recall that the isomorphism $I'_Q(e_Q(\sigma)) \simeq e_{Q'}(\sigma) \otimes_{(\mathcal{H}_{Q'}^+, j_{Q'}^+)} \mathcal{H}$ is given by $\varphi \mapsto \sum_{w \in Q' W_0} \varphi(T_{n_{w^{-1} w_G w_Q}}^*) \otimes T_{n_w}$ [Abe16, Lemma 2.22]. Let $\varphi' \in I'_{Q_1}(e_{Q_1}(\sigma))$ be the image of φ . Then the image of φ' in $e_{Q'_1}(\sigma) \otimes_{(\mathcal{H}_{Q'_1}^+, j_{Q'_1}^+)} \mathcal{H}$ is

$$\begin{aligned} & \sum_{w \in Q'_1 W_0} \varphi'(T_{n_{w^{-1} w_G w_{Q_1}}}^*) \otimes T_{n_w} \\ &= (-1)^{\ell(w_{Q_1} w_Q)} \sum_{w \in Q'_1 W_0} \varphi(T_{n_{w^{-1} w_G w_{Q_1}}}^* T_{n_{w_{Q_1} w_Q}}^*) \otimes T_{n_w}. \end{aligned}$$

Since $w_{Q_1} w_Q \in W_{0, Q_1}$ and $w^{-1} w_G w_{Q_1} \in W_0^{Q_1}$ [Abe16, Lemma 2.22], we have $\ell(w^{-1} w_G w_{Q_1}) + \ell(w_{Q_1} w_Q) = \ell(w^{-1} w_G w_Q)$. Hence we have $T_{n_{w^{-1} w_G w_{Q_1}}}^* T_{n_{w_{Q_1} w_Q}}^* = T_{n_{w^{-1} w_G w_Q}}^*$. Therefore

$$\sum_{w \in Q'_1 W_0} \varphi'(T_{n_{w^{-1} w_G w_{Q_1}}}^*) \otimes T_{n_w} = (-1)^{\ell(w_{Q_1} w_Q)} \sum_{w \in Q'_1 W_0} \varphi(T_{n_{w^{-1} w_G w_Q}}^*) \otimes T_{n_w}.$$

Let P_2 be a parabolic subgroup corresponding to $\Delta \setminus \Delta_P$. Let $w \in Q' W_0$ but $w \notin Q'_1 W_0$. Then there exists a simple reflection $s \in W_{0, Q'_1}$ such that $sw < w$.

Since $w \in {}^{Q'}W_0 \subset W_{0,P_2}$, we have $s \in S_{0,P_2}$. Hence for any $x \in e_{Q'_1}(\sigma)$, we have $x \otimes T_{n_w} = x \otimes T_{n_s}T_{n_{sw}} = xe_{Q'_1}(\sigma)(T_{n_s}^{Q'_1}) \otimes T_{n_{sw}} = 0$ since $e_{Q'_1}(\sigma)(T_{n_s}^{Q'_1}) = 0$. Hence

$$\begin{aligned} & (-1)^{\ell(w_{Q_1}w_Q)} \sum_{w \in {}^{Q'_1}W_0} \varphi(T_{n_w^{-1}w_Gw_Q}^*) \otimes T_{n_w} \\ &= (-1)^{\ell(w_{Q_1}w_Q)} \sum_{w \in {}^{Q'}W_0} \varphi(T_{n_w^{-1}w_Gw_Q}^*) \otimes T_{n_w}. \end{aligned}$$

Therefore the homomorphism

$$e_{Q'}(\sigma) \otimes_{(\mathcal{H}_{Q'_1}^+, j_{Q'_1}^+)} \mathcal{H} \rightarrow e_{Q'_1}(\sigma) \otimes_{(\mathcal{H}_{Q'_1}^+, j_{Q'_1}^+)} \mathcal{H}$$

is given by $x \otimes X \mapsto (-1)^{\ell(w_{Q_1}w_Q)} x \otimes X$. (Here we identify $x \in e_{Q'}(\sigma)$ with $x \in e_{Q'_1}(\sigma)$.)

The isomorphism

$$(e_{Q'}(\sigma) \otimes_{(\mathcal{H}_{Q'_1}^+, j_{Q'_1}^+)} \mathcal{H})^* \simeq \text{Hom}_{(\mathcal{H}_{Q'_1}^-, j_{Q'_1}^{-*})}(\mathcal{H}, e_Q(\sigma)^*) = I_Q(e_Q(\sigma)^*)$$

is given by $f \mapsto (X \mapsto (x \mapsto f(x \otimes \zeta(X))))$ and the opposite is given by $f' \mapsto ((x \otimes X) \mapsto f'(\zeta(X))(x))$. (Here we identify $e_Q(\sigma)^*$ with $e_Q(\sigma^*)$.) Hence the maps

$$I_{Q_1}(e_{Q_1}(\sigma)^*) \simeq (e_{Q'_1}(\sigma) \otimes_{(\mathcal{H}_{Q'_1}^+, j_{Q'_1}^+)} \mathcal{H})^* \rightarrow (e_{Q'}(\sigma) \otimes_{(\mathcal{H}_{Q'_1}^+, j_{Q'_1}^+)} \mathcal{H})^* \simeq I_Q(e_Q(\sigma)^*)$$

send $f \in I_{Q_1}'(e_{Q_1}(\sigma)^*)$ to

$$\begin{aligned} & (x \otimes X) \mapsto f(\zeta(X))(x) \in (e_{Q'_1}(\sigma) \otimes_{(\mathcal{H}_{Q'_1}^+, j_{Q'_1}^+)} \mathcal{H})^*, \\ & (x \otimes X) \mapsto (-1)^{\ell(w_{Q_1}w_Q)} f(\zeta(X))(x) \in (e_{Q'}(\sigma) \otimes_{(\mathcal{H}_{Q'_1}^+, j_{Q'_1}^+)} \mathcal{H})^*, \\ & X \mapsto (x \mapsto (-1)^{\ell(w_{Q_1}w_Q)} f(X)(x)) \in I_Q(e_Q(\sigma)^*). \end{aligned}$$

Namely, it is equal to the the natural embedding times $(-1)^{\ell(w_{Q_1}w_Q)}$. $\square$

4.3. Supersingular modules. Assume that C is a field.

Theorem 4.8. *Let (χ, J, V) be as in subsection 2.12. Then we have $\pi_{\chi, J, V}^* \simeq \pi_{\chi^{-1}, J, V^*}$.*

Proof. Let Ξ be a character of $\mathcal{H}_{\text{aff}}$ determined by (χ, J) . By the proof of [Vig15a, Proposition 6.17], $\Xi \otimes V \subset \pi_{\chi, J, V}|_{\mathcal{H}_{\Xi}}$ is a direct summand. Hence $(\Xi \otimes V)^* \subset (\pi_{\chi, J, V})^*|_{\mathcal{H}_{\Xi}}$. Since Ξ and V are finite-dimensional, we have $(\Xi \otimes V)^* = \Xi^* \otimes V^*$. Therefore we have a non-zero homomorphism $(\Xi^* \otimes V^*) \otimes_{\mathcal{H}_{\Xi}} \mathcal{H} \rightarrow \pi_{\chi, J, V}^*$. The restriction of V^* to $Z_{\kappa} \cap W_{\text{aff}, P}(1)$ is the direct sum of $\chi^* = \chi^{-1}$ since $V|_{Z_{\kappa} \cap W_{\text{aff}, P}(1)}$ is a direct sum of χ . For $s \in S_{\text{aff}, \chi} = S_{\text{aff}, \chi^{-1}}$, $\Xi^*(T_s) = \Xi(\zeta(T_s)) = \Xi(T_{\tilde{s}^{-1}})$ where $\tilde{s}$ is a lift of s . This is 0 or $\chi(c_{s^{-1}})$ and 0 if and only if $s \in J$. Hence the subset of $S_{\text{aff}, \chi^{-1}} = S_{\text{aff}, \chi}$ attached to Ξ^* is J . Therefore $(\Xi^* \otimes V^*) \otimes_{\mathcal{H}_{\Xi}} \mathcal{H} = \pi_{\chi^{-1}, J, V^*}$. Hence we get a non-zero homomorphism $\pi_{\chi^{-1}, J, V^*} \rightarrow \pi_{\chi, J, V}^*$. Since this is a non-zero homomorphism between simple modules, this is an isomorphism. $\square$

4.4. Simple modules. Assume that C is a field. Combining Proposition 4.2, 4.5 and Theorem 4.8, we get the following theorem.

Theorem 4.9. *Set $P' = n_{w_Gw_P} P^{\text{op}} n_{w_Gw_P}^{-1}$ and $Q' = n_{w_Gw_Q} Q^{\text{op}} n_{w_Gw_Q}^{-1}$. Let (χ', J', V') be a triple for $\mathcal{H}_{P'}$ defined by the pull-back of the triple (χ^{-1}, J, V^*) by $n_{w_Gw_P}$. Then we have $I(P; \chi, J, V; Q)^* = I(P'; \chi', J', V'; Q')$.*

We use the following lemma.

Lemma 4.10. *Let P be a parabolic subgroup and σ an $\mathcal{H}_P$ -module. Then we have $P(n_{w_G w_P} \sigma) = n_{w_G w_P(\sigma)} P(\sigma)^{\text{op}} n_{w_G w_P(\sigma)}^{-1}$.*

Remark 4.11. By [Abe16, Lemma 2.27], we have $n_{w_G w_P(\sigma)} \sigma = n_{w_G w_P} \sigma$.

Proof. Let P_α be the parabolic subgroup corresponding to $\Delta_P \cup \{\alpha\}$ where $\alpha \in \Delta \setminus \Delta_P$. Set $n = n_{w_G w_P}$ and $P' = n_{w_G w_P} P^{\text{op}} n_{w_G w_P}^{-1}$. Let $\alpha \in \Delta(n\sigma) \setminus \Delta_{P'}$ and we prove that $\alpha \in -w_G(\Delta(\sigma) \setminus \Delta_P)$. Since $\Delta_{P'} = -w_G(\Delta_P)$, $\langle \alpha, \Delta_{P'} \rangle = 0$ implies $\langle -w_G(\alpha), \Delta_P \rangle = 0$. If for any $\lambda \in \Lambda(1) \cap W_{\text{aff}, P_\alpha}(1)$ satisfies $(n\sigma)(T_\lambda^{P'}) = 1$, then we have $\sigma(T_{n^{-1}\lambda n}^P) = 1$. Hence $\sigma(T_\lambda^P) = 1$ for any $\lambda \in n^{-1}(\Lambda(1) \cap W_{\text{aff}, P_\alpha}(1))n = \Lambda(1) \cap W_{\text{aff}, P_{-(w_G w_P)^{-1}(\alpha)}}(1)$. Since $\langle -w_G(\alpha), \Delta_P \rangle = 0$, we have $-(w_G w_P)^{-1}(\alpha) = -w_P w_G(\alpha) = -w_G(\alpha)$. Hence $-w_G(\alpha) \in \Delta(\sigma)$. $\square$

Proof of Theorem 4.9. Set $\sigma = \pi_{\chi, J, V}$ and $P(\sigma)' = n_{w_G w_P(\sigma)} P(\sigma)^{\text{op}} n_{w_G w_P(\sigma)}^{-1}$. We have

$$\begin{aligned} I(P; \chi, J, V; Q)^* &= I_{P(\sigma)}(\text{St}_Q^{P(\sigma)} \sigma)^* \\ &\simeq I_{P(\sigma)'}(n_{w_G w_P(\sigma)} (\text{St}_Q^{P(\sigma)} \sigma)^*) \\ &\simeq I_{P(\sigma)'}(n_{w_G w_P(\sigma)} (\text{St}_{n_{w_P(\sigma)w_Q} Q^{\text{op}} n_{w_P(\sigma)w_Q}^{-1}}^{P(\sigma)} \sigma^*)). \end{aligned}$$

The adjoint action of $n_{w_G w_P(\sigma)}$ induces an isomorphism $\mathcal{H}_{P(\sigma)} \simeq \mathcal{H}_{P(\sigma)'}$. For a parabolic subgroup Q_1 between $P(\sigma)$ and P , let Q_2 be a parabolic subgroup corresponding to $w_G w_P(\sigma)(\Delta_{Q_1})$. Then the adjoint action of $n_{w_G w_P(\sigma)}$ induces an isomorphism $\mathcal{H}_{Q_1} \simeq \mathcal{H}_{Q_2}$ and sends $\mathcal{H}_{Q_1}^{P(\sigma)-}$ to $\mathcal{H}_{Q_2}^{P(\sigma)'}-$. Moreover, it is compatible with homomorphisms $j_{Q_1}^{P(\sigma)-*}$ and $j_{Q_2}^{P(\sigma)'}-*$. Hence we have $n_{w_G w_P(\sigma)} I_{Q_1}^{P(\sigma)}(e_{Q_1}(\sigma)) \simeq I_{Q_2}^{P(\sigma)'}(n_{w_G w_P(\sigma)} e_{Q_1}(\sigma))$.

Since $\Delta_{P(\sigma)} \setminus \Delta_P$ is orthogonal to Δ_P , $w_{P(\sigma)} w_P(\Delta_P) = \Delta_P$. Hence we have $w_G w_P(\sigma)(\Delta_P) = w_G w_P(\Delta_P) = \Delta_{P'}$. Therefore, from the above compatibility of negative algebras and homomorphisms, $n_{w_G w_P(\sigma)} e_{Q_1}(\sigma) = e_{Q_2}(n_{w_G w_P(\sigma)} \sigma)$ where e_{Q_2} is the extension from P' to Q_2 . Therefore, combining the formula in the above paragraph, we get $n_{w_G w_P(\sigma)} \text{St}_{Q_1}^{P(\sigma)}(\sigma) \simeq \text{St}_{Q_2}^{P(\sigma)'}(n_{w_G w_P(\sigma)} \sigma)$.

Now set $Q_1 = n_{w_P(\sigma)w_Q} Q^{\text{op}} n_{w_P(\sigma)w_Q}$. Then $\Delta_{Q_1} = w_{P(\sigma)} w_Q(\Delta_Q)$. Hence $\Delta_{Q_2} = w_G w_Q(\Delta_Q) = \Delta_{Q'}$. Therefore we have

$$I(P; \chi, J, V; Q)^* \simeq I_{P(\sigma)'}(\text{St}_{Q'}^{P(\sigma)'} n_{w_G w_P(\sigma)} \sigma^*).$$

By [Abe16, Lemma 2.27], $\sigma^* = n_{w_P(\sigma)w_P} \sigma^*$. Hence we get

$$I(P, \sigma, Q)^* \simeq I_{P(\sigma)'}(\text{St}_{Q'}^{P(\sigma)'} n_{w_G w_P} \sigma^*).$$

$\square$

REFERENCES

- [Abe] N. Abe, *Modulo p parabolic induction of pro- p -Iwahori Hecke algebra*, J. Reine Angew. Math., DOI:10.1515/crelle-2016-0043.
- [Abe16] N. Abe, *Parabolic inductions for pro- p -Iwahori Hecke algebras*, arXiv:1612.01312.
- [AHHV17] N. Abe, G. Henniart, F. Herzig, and M.-F. Vignéras, *A classification of irreducible admissible mod p representations of p -adic reductive groups*, J. Amer. Math. Soc. **30** (2017), no. 2, 495–559.

- [AHV17] N. Abe, G. Henniart, and M.-F. Vignéras, *On pro- p -Iwahori invariants of R -representations of reductive p -adic groups*, arXiv:1703.10384.
- [Deo77] V. V. Deodhar, *Some characterizations of Bruhat ordering on a Coxeter group and determination of the relative Möbius function*, Invent. Math. **39** (1977), no. 2, 187–198.
- [Oll14] R. Ollivier, *Compatibility between Satake and Bernstein isomorphisms in characteristic p* , Algebra Number Theory **8** (2014), no. 5, 1071–1111.
- [Vig15a] M.-F. Vignéras, *The pro- p -Iwahori Hecke algebra of a p -adic group III*, J. Inst. Math. Jussieu (2015), 1–38.
- [Vig15b] M.-F. Vignéras, *The pro- p Iwahori Hecke algebra of a reductive p -adic group, V (parabolic induction)*, Pacific J. Math. **279** (2015), no. 1-2, 499–529.
- [Vig16] M.-F. Vignéras, *The pro- p -Iwahori Hecke algebra of a reductive p -adic group I*, Compos. Math. **152** (2016), no. 4, 693–753.

DEPARTMENT OF MATHEMATICS, HOKKAIDO UNIVERSITY, KITA 10, NISHI 8, KITA-KU, SAPPORO, HOKKAIDO, 060-0810, JAPAN

E-mail address: abenori@math.sci.hokudai.ac.jp